

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 033 407 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(43) Date of publication:
06.09.2000 Bulletin 2000/36

(21) Application number: 99944770.9

(22) Date of filing: 22.09.1999

(51) Int. Cl.⁷: **C12N 15/67**, C12N 15/52,
C12P 13/04, C12P 13/14,
C12P 19/38, C12N 9/02,
C12N 9/10, C12N 1/21

(86) International application number:
PCT/JP99/05175

(87) International publication number:
WO 00/18935 (06.04.2000 Gazette 2000/14)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: 25.09.1998 JP 27178698
25.09.1998 JP 27178798

(71) Applicant: Ajinomoto Co., Inc.
Tokyo 104-0031 (JP)

(72) Inventors:
• ASAKURA, Yoko,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)
• NAKAMURA, Jun,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)
• KANNO, Sohei,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)
• SUGA, Mikiko,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)

• KIMURA, Eiichiro,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)
• ITO, Hisao,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)
• MATSUI, Kazuhiko,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)
• OHSUMI, Tsuyoshi,
c/o Ajinomoto Co., Inc.
Tokyo 104-0031 (JP)
• NAKAMATSU, Tsuyoshi,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)
• KURAHASHI, Osamu,
c/o Ajinomoto Co., Inc.
Kawasaki-shi, Kanagawa 210-0801 (JP)

(74) Representative: HOFFMANN - EITLE
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)


(54) **PROCESS FOR CONSTRUCTING AMINO ACID-PRODUCING BACTERIUM AND PROCESS FOR PRODUCING AMINO ACID BY FERMENTATION METHOD WITH THE USE OF THE THUS CONSTRUCTED AMINO ACID-PRODUCING BACTERIUM**

(57) A method of producing coryneform bacteria having an improved amino acid- or nucleic acid-productivity comprises the steps of introducing a mutation in a promoter sequence of amino acid- or nucleic acid-biosynthesizing genes on a chromosome of a coryneform bacterium to make it close to a consensus sequence or introducing a change in a promoter sequence of amino acid- or nucleic acid-biosynthesizing genes on a chromosome of a coryneform bacterium by gene recombination to make it close to a consensus sequence, to obtain mutants of the coryneform amino acid- or nucleic acid-producing microorganism, culturing the mutants and select a mutant capable of producing the intended

amino acid or nucleic acid in a large amount. This method can construct a mutant capable of suitably enriching or controlling the expression of an intended gene without using a plasmid and also capable of producing amino acids in a high yield, by the recombination or mutation.

EP 1 033 407 A1

FIG. 1



Description

Background of the Invention

5 [0001] The present invention relates to a method of constructing a mutant strain capable of producing amino acids in a high yield, and a method of producing L-amino acids by the fermentation with the mutant.

[0002] Methods of constructing mutant strains usable for the production of amino acids by the fermentation can be roughly classified into two methods. One of them comprises introducing random mutations into DNA with a chemical mutagen, and the other comprises the genetic recombination. In the latter method, a strain having an improved capacity
10 of producing an intended substance can be developed by enhancing a gene on a metabolic pathway relating to the biosynthesis of an intended substance, or by weakening a gene of an enzyme relating to the destruction. In this connection, for enhancing an intended gene, a plasmid capable of autonomously replicating independently from the chromosome in a cell has been mainly used.

[0003] However, the method of enhancing the intended gene with a plasmid has problems. In particular, the degree
15 of enrichment of the intended gene is variable depending on the number of copies of the plasmid itself. Therefore, for some kinds of intended genes, the copies are often too many in number and, as a result, the expression becomes excessive, the growth is seriously inhibited or the capacity of producing the intended substance is lowered. In such a case, although the degree of the enhancement of the intended gene can be lowered by using a plasmid of a small number of the copies, the variety of the plasmid is limited in many cases, and the intended control of the expression
20 level of the intended gene is impossible.

[0004] Another problem is that since the replication of the plasmid is often unstable, the plasmid is eliminated.

[0005] For example, Japanese Patent Unexamined Published Application (hereinafter referred to as "J. P. KOKAI") No.61-268185 discloses a recombinant DNA comprising a DNA fragment containing a glutamate dehydrogenase (GDH)-producing gene (glutamate dehydrogenase gene) derived from a glutamate-producing coryneform bacterium,
25 and a DNA fragment (plasmid) containing a gene necessary for the autonomous replication in the cell. It is also disclosed therein that by introducing the recombinant DNA into a cell, a GDH-enriching strain can be grown to improve the production of substances (such as amino acids and proteins) with microorganisms.

[0006] On the other hand, in Japanese Patent No. 2,520,895, the above described recombinant DNA is introduced into Corynebacterium to obtain a strain having the improved enzymatic activity, and L-glutamic acid is produced by the
30 fermentation with the strain. However, the production and yield of L-glutamic acid were yet unsatisfactory. Thus, it is demanded to further improve the productivity of L-glutamic acid. It is reported that the demand had been attained by introducing a recombinant DNA comprising two kinds of genes, i.e. a glutamate dehydrogenase-producing gene derived from a glutamate-producing coryneform bacterium, and an isocitrate dehydrogenase (ICDH) gene, into a glutamate-producing coryneform bacterium.

[0007] Further, JP Kokai No.6-502548 discloses an expression system and a secretion system of Corynebacterium comprising a Corynebacterium strain and a secretory cassette comprising the first functional DNA sequence for the expression in the strain, the second DNA sequence encoding for amino acids, polypeptides and/or proteins and the third DNA sequence inserted between the first DNA sequence and the second DNA sequence, wherein the third DNA sequence encodes the protein element selected from PS1 and PS2 which guarantee the secretion of the amino acids,
40 polypeptides and/or proteins. Specifically, the secretion of polypeptides is disclosed therein and in particular, NTG mutagenesis was conducted with Corynebacterium and a mutant resistant to 4-fluoroglutamate (4FG) which is an analogue to glutamate is selected and subjected to the transformation with PCGL141. It is described therein that a strain having an enhanced expression of GDH can be obtained from the analogue resistant bacteria. It is also described therein that a mutation was observed in nucleotide sequence No.251 to No.266 of GDH promoter.

Disclosure of the Invention

[0008] The object of the present invention is to provide a method of constructing a mutant capable of suitably enhancing or controlling the expression of an intended gene without using a plasmid and also capable of producing
50 amino acids in a high yield, by gene recombination or mutation.

[0009] Another object of the present invention is to provide a promoter for GDH capable of imparting a capability of producing glutamic acid in a high yield to a Corynebacterium strain without seriously increasing the amount of by-produced aspartic acid and alanine.

[0010] Still another object of the present invention is to provide a GDH gene having a sequence of the above-described promoter for GDH.
55

[0011] A further object of the present invention is to provide a Corynebacterium strain having the above-described gene and capable of producing L-glutamic acid.

[0012] A further object of the present invention is to provide a method of producing amino acids by fermentation

wherein amino acid-producing microorganism thus constructed is used.

[0013] A further object of the present invention is to provide a fermentation method of producing glutamic acid at a low cost by increasing the yield of glutamic acid by using a glutamic acid-producing coryneform bacterium.

[0014] The present invention has been completed on the basis of a finding that the above-described problems can be efficiently solved by variously modifying the promoter of amino acid-biosynthesizing genes on a chromosome to control the amount of the expression of the intended genes. Particularly, the invention has been completed on the basis of a finding that the above-described problem can be efficiently solved by introducing a specific mutation into -35 region or -10 region which is a specific region of the promoter.

[0015] Namely, the present invention provides a method of producing coryneform bacteria having an improved amino acid- or nucleic acid-productivity, which comprises the steps of introducing a mutations in a promoter sequence of amino acid- or nucleic acid-biosynthesizing genes on a chromosome of a coryneform bacteria to make it close to a consensus sequence or introducing a change in a promoter sequence of amino acid- or nucleic acid-biosynthesizing genes on a chromosome of Coryneform bacteria by gene recombination to make it close to a consensus sequence, to obtain mutants of the coryneform amino acid- or nucleic acid-producing microorganism, culturing the mutants, and selecting a mutant capable of producing the intended amino acid or nucleic acid in a large amount.

[0016] The present invention also provides a promoter for glutamate dehydrogenase (GDH)-producing gene, which has the sequence of (i) at least one DNA sequence selected from the group consisting of CCGTCA, TTGTCA, TTGACA and TTGCCA in -35 region, (ii) TATAAT sequence or the same TATAAT sequence but in which the base of ATAAT is replaced with another base in -10 region, or (iii) a combination of (i) and (ii), wherein the sequence does not inhibit the promoter function.

[0017] The present invention also provides a glutamate dehydrogenase-producing gene having the above-described promoter.

[0018] The present invention also provides a coryneform L-glutamate-producing microorganism having the above-described gene.

[0019] The present invention also provides a process for producing an amino acid by the fermentation, which comprises the steps of culturing a coryneform bacterium constructed by the above-described method and having an improved amino acid-producing capacity in a medium to form and also to accumulate the intended amino acid in the medium, and collecting the amino acid from the medium.

[0020] The present invention also provides a process for producing L-glutamic acid by the fermentation, which comprises the steps of culturing a coryneform L-glutamic acid-producing microorganism resistant to 4-fluoroglutamic acid in a liquid medium to form and also to accumulate L-glutamic acid in the medium, and collecting L-glutamic acid from the medium.

Brief Description of the Drawings

[0021]

Fig.1 show a flow of construction of GDH gene having a mutant promoter.

Fig.2 show a flow of construction of CS gene having a mutant promoter.

Fig.3 show a flow of construction of shuttle vector carrying lacZ as a reporter gene.

Best Mode for Carrying out the Invention

[0022] The term "coryneform glutamic acid producing microorganism" as used herein includes also bacteria which were classified to be the genus *Brevibacterium* before but now integrated into the genus *Corynebacterium* [Int. J. Syst. Bacteriol., 41, 255 (1981)] and also bacteria of the genus *Brevibacterium* which are very close to those of the genus *Corynebacterium*. Therefore, the mutants used in the present invention can be derived from the coryneform glutamic acid-producing bacteria of the genus *Brevibacterium* or *Corynebacterium* shown below. Bacteria of the genus *Corynebacterium* and those of the genus *Brevibacterium* will be collectively referred to as "coryneform bacteria" so far as they do not concern the glutamic acid productivity.

<i>Corynebacterium acetoacidophilum</i>	ATCC13870
<i>Corynebacterium acetoglutamicum</i>	ATCC15806
<i>Corynebacterium callunae</i>	ATCC15991
<i>Corynebacterium glutamicum</i>	ATCC13032
<i>Brevibacterium divaricatum</i>	ATCC14020
<i>Brevibacterium lactofermentum</i>	ATCC13869
<i>Corynebacterium lilium</i>	ATCC15990

<i>Brevibacterium flavum</i>	ATCC14067
<i>Corynebacterium melassecola</i>	ATCC17965
<i>Brevibacterium saccharolyticum</i>	ATCC14066
<i>Brevibacterium immariophilum</i>	ATCC14068
5 <i>Brevibacterium roseum</i>	ATCC13825
<i>Brevibacterium thioenitalis</i>	ATCC19240
<i>Microbacterium ammoniaphilum</i>	ATCC15354
<i>Corynebacterium thermoaminogenes</i>	AJ12310(FERM 9246)

10 [0023] The amino acids to be produced are not particularly limited so far as the genes concerning the biosynthesis and promoters thereof have been elucidated. Examples of effective enzymes concerning the biosynthesis include GDH, citrate synthase (CS), isocitrate synthase (ICDH), pyruvate dehydrogenase (PDH) and aconitase (ACO) for glutamic acid fermentation.

15 [0024] Enzymes for lysine fermentation including biosynthesis enzymes such as aspartate kinase (AK), dihydrodipicolinate synthase, dihydrodipicolinate reductase, diaminopimelate dehydrogenase and diaminopimelate decarboxylase are also effective. Lysine eccrisis protein (lysE gene) concerning the membrane eccrisis of lysine is also effective.

[0025] Effective enzymes for arginine fermentation include N-acetylglutamate synthase, N-acetylglutamate kinase, N-acetylglutamyl phosphate reductase, acetylornithine aminotransferase, N-acetylornithinase, ornithine carbamyltransferase, argininosuccinate synthase, and argininosuccinase. arginine is formed by the reaction catalyzed by these enzymes. These enzymes are effective. These enzymes are coded by enzymes argA, argB, argC, argD, aegE, argF, argG and argH, relatively.

[0026] Effective enzymes for serine fermentation includes 3-phosphoglyceric acid dehydrogenase, phosphoserine trans-aminase, phosphoserine phosphatase and the like.

25 [0027] Effective enzymes for phenylalanine fermentation include bio-synthesizing enzymes such as deoxyarabinohepturonic phosphate synthetase, 3-dehydrokinate synthetase, 3-dehydrokinic acid dehydratase, shikimic acid dehydrogenase, shikimic kinase, 5-enol pyruvylshikimic acid-3-phosphate synthetase, chorismic acid synthetic enzyme, chorismate synthetase, chorismate mutase, prephenate dehydratase, and the like. Sugar metabolic enzymes such as transketorase, transaldolase, phosphoenolpyruvic acid synthetic enzyme are also effective.

[0028] Effective enzymes for tryptophan fermentation include enzymes belonging to tryptophan operon, in addition to various enzymes effective in the above-mentioned phenylalanine fermentation and various enzymes effective in the above-mentioned serine fermentation.

[0029] Effective enzymes for proline fermentation include γ -glutamylkinase, γ -glutamylcemicialdehyde dehydrogenase, pyrroline-5-carboxylate reductase, in addition to various enzymes effective in the above-mentioned glutamic acid fermentation.

35 [0030] Effective enzymes for glutamine fermentation include glutamine synthetase, in addition to various enzymes effective in the above-mentioned glutamic acid fermentation.

[0031] In the inosine production, it is considered to be useful to enhance the expression of 5-phosphoribosyl 1-diphosphate synthetase, 5-phosphoribosyl 1-diphosphate aminotransferase, phosphoribosylaminoimidazole-carboxamide formyltransferase and the like.

40 [0032] In the guanosine production, it is considered to be useful to enhance the expression of 5'-inosinic acid dehydrogenase and 5'-xanthylic acid aminase, addition to 5-phosphoribosyl 1-diphosphate synthetase, 5-phosphoribosyl 1-diphosphate aminotransferase, phosphoribosylaminoimidazolecarboxamide formyltransferase and the like.

[0033] In the adenosine production, it is considered to be useful to enhance the expression of adenosuccinate synthase, in addition to 5-phosphoribosyl 1-diphosphoric acid synthetic enzyme, 5-phosphoribosyl 1-diphosphoric acid aminotransferase, phosphoribosylaminoimidazole-carboxamide formyltransferase and the like.

45 [0034] In the nucleotide production, it is considered to be useful to enhance the expression of phosphoribosyl transferase, inosine kinase, guanosine kinase and adenosine kinase.

[0035] In the present invention, a mutant of a coryneform amino acid-producing bacterium is obtained by, introducing a mutation in a promoter sequence of desired amino acid-biosynthesizing genes on a chromosome of a coryneform amino acid-producing bacterium, such as the above-described promoter sequence for GDH, to make it close to a consensus sequence with a chemical or by introducing the mutation by the genetic recombination to obtain a mutant of the coryneform amino acid-producing microorganism.

55 [0036] The term "consensus sequence" is a sequence which appears most frequently in various promoter sequences. Such consensus sequences include, for example, those of *E. coli* and *Bacillus subtilis*. The consensus sequence of *E. coli* is described in Diane K. Hawley and William R. McClure Nuc. Acid. Res. 11:2237-2255(1983), and that of *B. subtilis* is described in Charles et al. Mol. Gen. Genet 186:339-346(1982).

[0037] The mutation may be caused in either only one promoter sequence such as that for GDH or two or more promoter sequences such as those for GDH, citrate synthase (citrate-synthesizing enzyme) (CS) and isocitrate synthase

(isocitrate-synthesizing enzyme) (ICDH).

[0038] In the present invention, the mutant thus obtained is cultured to obtain the mutant capable of producing a large amount of an intended amino acid.

[0039] It was already elucidated that in the fermentation of glutamic acid, GDH derived from a coryneform glutamate-producing microorganism has its own promoter sequence in upstream region thereof [Sahm et al., Molecular Microbiology (1992), 6, 317-326].

[0040] For example, the promoter for GDH of the present invention, GDH gene having the promoter sequence for GDH and L-glutamate-producing *Corynebacterium* strain having this gene can be obtained by, for example, the following methods:

[0041] Namely, the strain is subjected to a mutagenesis treatment such as the irradiation with UV, X-rays or radiation, or treatment with a mutagen to obtain a strain resistant to 4-fluoroglutamic acid on an agar plate culture medium containing 4-fluoroglutamic acid. Namely, the mutagenized cells are spread on agar plates culture medium containing 4-fluoroglutamic acid in such a concentration that it inhibits the growth of the parent, and the mutant thus grown is separated.

[0042] Further, the promoter sequence of GDH genes can be replaced with variously modified sequences by site directed mutagenesis, and the relationship between the respective sequences and GDH activity is examined so as to select the ones having a high L-glutamate-productivity.

[0043] It is particularly preferred in the present invention that the DNA sequence in -35 region of the promoter for GDH-producing gene is at least one DNA sequence selected from the group consisting of CGGTCA, TTGTCA, TTGACA and TTGCCA and/or the DNA sequence in -10 region of the promoter is TATAAT, or the bases of ATAAT in TATTAT sequence in -10 region is replaced with another base, while they do not inhibit the promoter function. The reason why the strain in which the bases of ATAAT in TATAAT sequence in -10 region is replaced with another base and the promoter function is not inhibited can be selected is as follows: Because a remarkable increase in the specific activity of GDH was observed by merely replacing the first "C" of CATAAT with "T" in wild type -10 sequence (refer to p6-4 in Table 1), it was considered that such a replacement with another base is possible.

[0044] The promoter sequence of GDH gene is described in, for example, the above-described Sahm et al., Molecular Microbiology (1992), 6, 317-326. It is described therein as Seq ID No. 1. The sequence of GDH gene itself is also described in Sahm et al., Molecular Microbiology (1992), 6, 317-326 to be Seq ID No. 1.

[0045] Similarly, the mutation can be introduced in the promoter for citrate-synthesizing enzyme (CS) or isocitrate-synthesizing enzyme (ICDH).

[0046] Thus, the promoters for GDH are those having at least one DNA sequence in -10 region selected from the group consisting of CGGTCA, TTGTCA, TTGACA and TTGCCA in -35 region and/or or TATAAT sequence or the TATAAT sequence but in which the base of ATAAT is replaced with another base, wherein they do not inhibit the promoter function. Genes for producing glutamate dehydrogenase, which have the above-described promoter, are also provided.

[0047] The promoters for CS are Those having TTGACA sequence in -35 region and/or TATAAT sequence in -10 region, which do not inhibit the promoter function. CS genes having the above-described promoter are also provided.

[0048] Promoters for ICDH are those having TTGCCA or TTGACA sequence in the first or the second promoter in -35 region and/or TATAAT sequence in the first or the second promoter in -10 region which do not inhibit the function of the promoter. The *icd* genes having the above-described promoter are also provided.

[0049] Promoters for PDH are those having TTGCCA sequence in -35 region and/or TATAAT sequence in -10 region, which do not inhibit the function of the promoter. PDH genes having the above-described promoter are also provided.

[0050] The present invention also provides coryneform L-glutamate-producing bacterium having the above-described genes.

[0051] The promoters for argininosuccinate synthase are those having at least one DNA sequence selected from the group consisting of TTGCCA, TTGCTA, and TTGTCA in -35 region and/or TATAAT sequence in -10 region, or the base of ATAAT in TATTAT sequence is replaced with another base, which do not inhibit the function of the promoter. Argininosuccinate synthase gene having the above-described promoter are also provided.

[0052] The present invention also provides coryneform arginine-producing bacterium having the above-described genes.

[0053] Amino acids can be obtained by culturing a coryneform bacterium of the present invention, which produces an amino acid, preferably L-glutamic acid, in a liquid culture medium to form and thereby to accumulate the intended amino acid, preferably L-glutamic acid, and collecting the amino acid from the culture medium.

[0054] The liquid culture medium used for cultivating the above-described strain of the bacterium in the present invention is an ordinary nutrition medium containing carbon sources, nitrogen sources, inorganic salts, growth factors, etc.

[0055] The carbon sources include carbohydrates such as glucose, fructose, sucrose, molasses and starch hydro-

lyzates; alcohols such as ethanol and glycerol; and organic acids such as acetic acid. The nitrogen sources include ammonium sulfate, ammonium nitrate, ammonium chloride, ammonium phosphate, ammonium acetate, ammonia, peptone, meat extract, yeast extract and corn steep liquor. When an auxotrophic mutant is used, the required substances are added to the medium as the reagents or natural substances containing them.

5 [0056] The coryneform bacteria usually produce L-glutamic acid under reduced biotin condition. Therefore, the amount of biotin in the medium is restricted or a substance inhibiting the effect of biotin such as a surfactant or penicillin is added.

[0057] The fermentation is preferably conducted by shaking the culture or agitating the culture with aeration while the pH of the culture liquid is kept in the range of 5 to 9 for 2 to 7 days. The pH is preferably controlled with urea, calcium carbonate, gaseous ammonia, ammonia water or the like. The culture temperature is preferably 24 to 37°C.

10 [0058] L-glutamic acid thus produced and accumulated in the culture liquid is collected by an ordinary method such as ion-exchange resin method or crystallization method. Specifically, L-glutamic acid is separated by the adsorption on an anion-exchange resin or by the neutralization crystallization.

[0059] According to the present invention, the intended amino acid can be obtained in a high yield by introducing a mutation into a promoter region of amino acid-biosynthesizing genes of a coryneform amino acid-producing bacterium to control the expression of the intended genes. In addition, since any elimination of the intended gene does not occur in the bacteria according to the present invention, contrary to the cases using plasmid, the intended amino acid can be stably obtained in a high yield. Thus, the industrial merit of the invention is great.

20 [0060] The present invention provides various promoters, particularly, promoters for GDH, capable of imparting a power of producing amino acids, particularly glutamic acid, in a high yield to Corynebacterium strains without increasing the amount of by-produced aspartic acid and alanine.

[0061] In the present invention, a coryneform L-glutamate-producing bacterium is mutagenized, a strain in which the mutation introduced in a promoter region of GDH gene and which is resistant to 4-fluoroglutamic acid is collected, and the strain is cultured to obtain glutamic acid in a high yield. Thus, the present invention is industrially very advantageous.

25 [0062] The following Examples will further illustrate the present invention.

Example 1: Production of mutant GDH promoter:

30 [0063] A mutant GDH promoter was prepared by site-directed mutagenesis method as follows:

(1) Preparation of GDH genes having various mutant promoters:

[0064] The wild type sequence in -35 region and -10 region of a promoter of GDH gene of a coryneform bacteria is shown in sequence 1. The promoter sequence of wild type has already been reported [Molecular Microbiology (1992), 6, 317-326].

[0065] The method of preparing a plasmid carrying GDH gene having a mutant promoter is as follows:

[0066] As shown in Fig. 1, a chromosomal gene of a wild type strain of a coryneform bacterium ATCC13869 prepared with "Bacterial Genome DNA purification kit" (Advanced Genetic Technologies Corp.) was used as the template for PCR. The gene amplification was conducted by PCR using upstream and downstream sequences of GDH gene. Both ends were blunt-ended. The product thus obtained was inserted in SmaI site of plasmid pHSG399 (a product of Takara Shuzo Co., Ltd.). Then a replication origin taken from plasmid pSAK4 having the replication origin capable of replicating in a coryneform bacterium was introduced into Sal I site of the plasmid to obtain plasmid pGDH. By this method, GDH genes having each above-described promoter sequence can be obtained by using a primer having each of the sequence of Seq ID No. 1 to Seq ID No. 6 shown in the Sequence Listing as the upstream primer for GDH gene, respectively. It was confirmed by sequencing the PCR amplified fragment that any mutation, other than the introduced mutation in the promoter sequence, was not occurred in the amplified fragment. pSAK4 is constructed as follows: previously obtained plasmid pHK4 (J. P. KOKAI No.5-7491) having an autonomous replication origin derived from plasmid pHM1519 [Agric. Biol. Chem., 48, 2901-1903 (1984)], which is capable of autonomously replicating in Corynebacterium microorganism, is digested with restriction enzymes BamHI and KpnI to obtain a DNA fragment having the replication origin. Then the fragment thus obtained is blunt-ended with DNA-Blunting Kit (Blunting kit of Takara Shuzo Co., Ltd.). After the ligation with SalI linker, the product thus obtained was inserted into Sal I site of pHSG299 (a product of Takara Shuzo Co., Ltd.) to obtain plasmid pSAK4.

(2) Comparison of the degrees of expression of GDH having each promoter sequence:

55 [0067] Each plasmid prepared as described above was introduced into wild type strain of coryneform bacterium ATCC13869 by electroporation method (refer to J. P. KOKAI No. 2-207791. For comparing the degrees of expression of

GDH for these strains, the specific activity of GDH was determined by the above-described method of Sahm et al. The results are shown in Table 1.

Table 1

Strain	Promoter sequence		Specific activity of GDH	Relative value
	-35	-10		
ATCC13869	TGGTCA	CATAAT	7.7	0.1
/pGDH	TGGTCA	CATAAT	82.7	1.0
/p6-2	CGGTCA	CATAAT	33.1	0.4
/p6-4	TGGTCA	TATAA.T	225.9	2.7
/p6-3	TTGACA	TATAAT	327.2	4.0
/p6-7	TTGCCA	TATAAT	407.0	4.9
/p6-8	TTGTCA	TATAAT	401.3	4.9

[0068] ATCC 13869/p6-2 through ATCC 13869p6-8/ corresponded to the sequences of Seq ID No. 2 through Seq ID No.6, respectively. These sequences were the same as the sequence No.1 (wild type) except that the underlined parts were changed as follows:

Sequence

[0069]

No. 1 5'-TTAATTCTTTGTGGTCAATATCTGCGACACTGC CATAATTTGAACGT-3'

2 CGGTCA CATAAT

3. TGGTCA TATAAT

4. TTGACA TATAAT

5. TTGCCA TATAAT

6. TTGTCA TATATT

[0070] These were those of synthetic linear doubled stranded DNA.

Example 2: Preparation of mutant strains:

(1) Preparation of mutant strains resistant to 4-fluoroglutamic acid:

[0071] AJ13029 is a mutant strain producing glutamic acid and disclosed in W096/06180. Although it does not produce glutamic acid at a culture temperature of 31.5°C, it produces glutamic acid even in the absence of a biotin-inhibitor when the culture temperature is shifted to 37°C. In this Example, *Brevibacterium lactofermentum* AJ13029 strain was used as the parent strain for preparing the mutant strains. As a matter of course, any of glutamic acid-producing strains other than AJ13029 can be used as a parent strain for preparing mutant strains resistant to 4-fluoroglutamic acid.

[0072] AJ13029 were cultured on a CM2B agar medium (Table 2) at 31.5°C for 24 hours to obtain the bacterial cells. The cells were treated with 250µg/ml aqueous solution of N-methyl-N'-nitro-N-nitrosoguanidine at 30°C for 30 minutes. Then a suspension of the cells having a survival rate of 1 % was spread on agar plates culture medium (Table 3) containing 4-fluoroglutamic acid (4FG). Colonies were formed after incubating the plate at 31.5°C for 20 to 30 hours. In this experiment, a slant medium containing 1 mg/ml of 4FG was prepared at first, and then a layer of the same

medium without 4FG was formed thereon horizontally. Thus, 4FG concentration gradient was obtained on the surface of the agar medium. When the plate was inoculated with the mutant cells obtained as described, a boundary line was formed at a border of the growing limit of the strain. Bacterial strains which formed colonies in a area containing 4FG of a concentration higher than that of the boundary line were taken. Thus, about 50 strains resistant to 4FG were obtained from about 10,000 mutagenized cells.

Table 2

CM2B agar medium	
Ingredient	Concentration
Polypeptone (Nippon Seiyaku Co.)	1.0 %
Yeast extract (Difco Co.)	1.0 %
NaCl	0.5 %
d-Biotin	10 µg/l
Agar	1.5%
(pH 7.2: adjusted with KOH)	

Table 3

Agar medium	
Component	Amount in one liter of water
Glucose	10 g
MgSO ₄ · 7H ₂ O	1g
FeSO ₄ · 7H ₂ O	0.01g
MnSO ₄ · 4-6H ₂ O	0.01 g
Thiamine hydrochloride	0.2 mg
d-Biotin	0.05 mg
(NH ₄) ₂ SO ₄	5 g
Na ₂ HPO ₄ · 12H ₂ O	7.1g
KH ₂ PO ₄	1.36g
Agar	15g

(2) Confirmation of capability of L-glutamic acid-production of 4FG-resistant mutant strains:

[0073] The capability of glutamic acid-production of about 50 mutant strains obtained in above (1) and parent AJ13029 strain were confirmed as described below.

[0074] AJ13029 and mutant strains were each cultured on CM2B agar medium at 31.5°C for 20 to 30 hours. A liquid medium having a composition shown as "medium A" in Table 4 was inoculated with the cells thus obtained, and the shaking culture was started at 31.5°C. About 22 hours after, the fresh medium was added so that the final concentration would be that of medium B shown in Table 4. The temperature was shifted to 37°C and then the culture was continued further for about 24 hours. After the completion of the culture, the culture was examined with a Biotic Analyzer (a product of Asahi Chemical Industry Co., Ltd.) to determine whether L-glutamic acid was produced or not. It was thus found that when the 50 strains were cultured, two strains having a yield of glutamic acid higher than that obtained from the parent strains and a high GDH activity were separated (strain A and strain B). GDH activity of each of them was determined to find that the specific GDH activity of both of them was increased (Table 5). The GDH activity was determined

by the method of E. R. Bormann et al. [Molecular Microbiol., 6, 317-326 (1996)]. By sequencing the GDH genes, it was identified that the mutation points were found only in the promoter region of GDH (Table 6).

Table 4

Ingredient	Medium A	Medium B
Glucose	3 g/dl	5 g/dl
KH ₂ PO ₄	0.14 g/dl	0.14 g/dl
MgSO ₄ · 7H ₂ O	0.04 g/dl	0.04 g/dl
FeSO ₄ · 7H ₂ O	0.001 g/dl	0.001 g/dl
MnSO ₄ · 4H ₂ O	0.001 g/dl	0.001 g/dl
(NH ₄) ₂ SO ₄	1.5 g/dl	2.5 g/dl
Soybean protein hydrolyzate solution	1.5 ml/dl	0.38 ml/dl
Thiamine hydrochloride	0.2 mg/l	0.2 mg/l
Biotin	0.3 mg/l	0.3 mg/l
Antifoaming agent	0.05ml/l	0.05ml/l
CaCO ₃	5 g/dl	5 g/dl
pH	7.0(adjusted with KOH)	

Table 5

Glutamic acid formation and GDH activity of mutant strains			
Strain	Glu(g/dl)	GDH specific activity	Relative value
AJ13029	2.6	7.7	1.0
FGR1	2.9	23.1	3.0
FGR2	3.0	25.9	3.4

Table 6. DNA sequences in GDH promoter region of mutant strains

Strain	GDH promoter sequence		
	-35		-10
AJ13029	TGGTCA	TTCTGTGCGACACTGC	CATAAT
FGR1	TGGTCA	TTCTGTGCGACACTGC	TATAAT

EGR2 TTGTCA T-CTGTGCGACACTGC TATAAT

Example 3 Introduction of mutation into CS gene promoter region of coryneform glutamate-producing bacterium:

[0075] In this Example, a strain having an enhanced promoter for the genes which codes glutamate dehydrogenase (GDH) and citrate-synthesizing enzyme (CS) was produced.

(1) Cloning of gltA gene:

[0076] The sequence of gltA gene of a coryneform bacterium, which codes citrate-synthesizing enzyme, has already been elucidated [Microbial. 140, 1817-1828 (1994)]. On the basis of this sequence, primers shown in Seq ID No. 7 and Seq ID No. 8 were synthesized. On the other hand, chromosomal DNA from *Brevibacterium lactofermentum* ATCC13869 was prepared using Bacterial Genome DNA Purification Kit (Advanced Genetic Technologies Corp.). Sterilized water was added to a mixture of 0.5 µg of the chromosomal DNA, 10 pmol of each of the oligonucleotides, 8 µl of dNTP mixture (2.5 mM each), 5 µl of 10xLa Taq Buffer (Takara Shuzo Co., Ltd.) and 2 U of La Taq (Takara Shuzo Co., Ltd.) to obtain 50 µl of PCR-reaction cocktail. The reaction cocktail was subjected to PCR. The PCR conditions were 30 cycles of denaturation at 94°C for 30 seconds, annealing at 55°C for 15 seconds and extension at 72°C for 3 seconds using Thermal Cycler TP240 (Takara Shuzo Co., Ltd.) to amplify about 3 Kbp of DNA fragments containing gltA gene and promoter thereof. The amplified fragments thus obtained were purified with SUPRECO2 (Takara Shuzo Co., Ltd.) and then blunt-ended. The blunting was conducted with Blunting Kit of Takara Shuzo Co., Ltd. The blunt-ended fragment was mixed with pHSG399 (Takara Shuzo Co., Ltd.) completely digested with SmaI to conduct the ligation. The ligation reaction was conducted with DNA Ligation Kit ver 2 (Takara Shuzo Co., Ltd.). After the completion of the ligation, the transformation was conducted with competent cells of *E. coli* JM109 (Takara Shuzo Co., Ltd.). The cells were spread on an L medium plates (comprising 10 g/l of bactotryptone, 5 g/l of bacto-yeast extract, 5 g/l of NaCl and 15 g/l of agar; pH 7.2) containing 10 µg/ml of IPTG (isopropyl-β-D-thiogalactopyranoside), 40 µg/ml of X-Gal (5-bromo-4-chloro-3-indolyl-β-D-galactoside) and 40 µg/ml of chloramphenicol. After culturing them overnight, white colonies were taken to obtain the transformed strains after single colony isolation.

[0077] From the transformed strains, plasmids were prepared by the alkali method (*Seibutsu Kogaku Jikken-sho* edited by Nippon Seibutsu Kogaku-kai and published by Baifukan, p. 105, 1992). Restriction enzyme maps were prepared, and the plasmid which has the same restriction map as the map shown in Fig. 2 was named "pHSG399CS".

(2) Introduction of mutations into gltA promoter:

[0078] Mutan-Super Express Km (Takara Shuzo Co., Ltd.) was used for the introduction of mutation into gltA promoter region. The method is specifically described below. pHSG399CS was completely digested with EcoRI and SmaI to obtain EcoRI-SmaI fragment containing gltA genes, which were ligated to the fragment obtained by complete digestion of pKF19kM (Takara Shuzo Co., Ltd.) with EcoRI and SmaI. After the completion of the ligation, the transformation was conducted with competent cells of *E. coli* JM109 (Takara Shuzo Co., Ltd.). The cells were spread on L medium plates containing 10 µg/ml of IPTG, 40 µg/ml of X-Gal and 25 µg/ml of kanamycin. After overnight incubation, white colonies were taken and transformants were obtained by single colony isolation. From the transformants, plasmids were prepared and the plasmid containing gltA gene was named pKF19CS.

[0079] PCR was conducted by using pKF19CS as the template and 5'-phosphorylated synthetic DNA shown in sequence of Seq ID No. 9, Seq ID No.10 and Seq ID No.11 together with the selection primer from Mutan super Express Km. The transformation was conducted with competent cells of *E. coli* MV1184 (Takara Shuzo Co., Ltd.) by using the PCR product. The cells were spread on L-medium plates containing 25 µg/ml of kanamycin. After overnight incubation, colonies were taken and the transformants were obtained after single colony isolation. From the transformants, plasmid DNA was prepared. The sequence of gltA promoter region was determined by Sanger method [J. Mol. Biol., 143,161 (1980)] using synthetic DNA having the sequence of Seq ID No. 12. Specifically, the sequence was determined with a Dye Terminator Sequencing Kit (Applied Biosystems) and analyzed by Genetic Analyzer ABI310 (Applied Biosystems). The plasmids in which gltA promoter region was replaced with the sequence shown in Table 7 were named pKF19CS1, pKF19CS2 and pKF19CS4, respectively.

Table 7

	-35 region	-10 region
pKF19CS	ATGGCT	TATAGC
pKF19CS1	ATGGCT	TATAAC
pKF19CS2	ATGGCT	TATAAT
pKF19CS4	TTGACA	TATAAT

(3) Construction of mutant *gltA* plasmid:

[0080] pKF19CS, pKF19CS1, pKF19CS2 and pKF19CS4 constructed in step (2) were completely digested with Sall and EcoRI (Takara Shuzo Co., Ltd.). On the other hand, plasmid pSFK6 (Japanese Patent Application No.11-69896) having a replication origin derived from plasmid pAM330 which can autonomously replicate in a coryneform bacterium [Japanese Patent Publication for Opposition Purpose (hereinafter referred to as "J. P. KOKAI") No. 58-67699] was completely digested with EcoRI and Sall. The obtained fragment was ligated with about the 2.5 kb fragment containing *gltA*. After the completion of the ligation, transformation was conducted with competent cells of *E. coli* JM109. The cells were spread on the L-medium plates containing 10 µg/ml of IPTG, 40 g/ml of X-Gal and 25 µg/ml of kanamycin. After overnight incubation, colonies were taken and the transformants were obtained after single colony isolation. From the transformants, plasmids were prepared. The plasmids containing *gltA* gene were named pSFKC, pSFKC1, pSFKC2 and pSFKC4, respectively.

(4) Determination of CS expression from mutant *gltA* plasmid in coryneform bacterium:

[0081] The plasmid constructed in above step (3) was introduced into *Brevibacterium lactofermentum* ATCC13869. Specifically, this treatment was conducted by electrical pulse method (J. P. KOKAI No. 2-07791). The transformants were selected at 31°C with CM2B medium plate (comprising 10 g/l of bactotryptone, 10 g/l of bacto yeast extract, 5 g/l of NaCl, 10 µg/l of biotin and 15 g/l of agar; pH 7.0) containing 25 µg/ml of kanamycin. After incubating for two days, colonies were taken and the transformants containing pSFKC, pSFKC1, pSFKC2 and pSFKC4 were named BLCS, BLCS1, BLCS2 and BLCS4, respectively, after single colony isolation. A medium having a composition shown in Table 8 was inoculated with the transformant. The culture was continued at 31°C and terminated before the glucose had been completely consumed. The culture liquid was centrifuged to separate the cells. The cells were washed with 50 mM tris buffer solution (pH 7.5) containing 200 mM of sodium glutamate and then suspended in the same buffer solution. After the sonication with UD-201 (TOMY) followed by the centrifugation (10,000g), the cells remaining unbroken were removed to obtain a crude enzyme solution. The activity of citrate synthase can be determined according to Methods Enzymol. 13, 3-11 (1969). Specifically, the crude enzyme solution was added to a reaction mixture containing 100 mM of TrisHCl (pH 8), 0.1 mM, of DTNB, 200 mM of sodium glutamate and 0.3 mM of acetyl CoA, and the background was determined as the increase in the absorbance at 412 nm at 30°C determined by Hitachi spectrophotometer U-3210. Further, oxaloacetic acid was added in such an amount that the final concentration thereof would be 0.5 mM. The increase in the absorbance at 412 nm was determined, from which the background value was deducted to determine the activity of the citrate synthase. The protein concentration in the crude enzyme solution was determined by Protein Assay (BIO-RAD). Bovine serum albumin was used as the standard protein. The results are shown in Table 9. It was confirmed that the citrate synthase activity of mutant *gltA* promoters was increased compared to wild-type *gltA* promoter.

Table 8

Ingredient	Concentration
Glucose	50 g/l
KH ₂ PO ₄	1 g/l
MnSO ₄ · 7H ₂ O	0.4 mg/l
FeSO ₄ · 7H ₂ O	10 mg/l
Soybean protein hydrolysates	20 ml/l

Table 8 (continued)

Ingredient	Concentration
Biotin	0.5 mg/l
Thiamine hydrochloride 2 mg/l	2 mg/l

Table 9

Strain	dABS/min/mg	Relative activity	Relative activity
Wild type 4	6.8	1.0	
BLCS00	38.8	5.7	1.0
BLCS01	57.1	8.4	1.21
BLCS02	92.5	13.6	1.9
BLCS04	239.5	35.2	4.8

(5) Introduction of mutant *gltA* gene into temperature-sensitive plasmid:

[0082] For integrating mutant *gltA* promoter sequences into a chromosome, a method is known wherein a plasmid of which replication in a coryneform bacterium is temperature sensitive is used (J. P. KOKAI No. 5-7491). PSFKT2 (Japanese Patent Application No. 11-81693) was used as the plasmid vector, the replication of which in a coryneform bacterium is temperature sensitive. pKFCS1, pKFCS2 and pKFCS3 completely digested with *SaiI* and *BstPI* and blunt-ended were used as the mutant *gltA* promoter sequences. They were ligated to pSFKT2 completely digested with *SmaI*. After the completion of the ligation, the transformation was conducted with competent cells of *E. coli* JM109 (Takara Shuzo Co., Ltd.). The cells were spread on the L-medium plates containing 10 μ g/ml of IPTG, 40 μ g/ml of X-Gal and 25 μ g/ml of kanamycin. After overnight incubation, white colonies were taken and the transformants were obtained after single colony isolation. From the transformants, plasmids were prepared. Temperature-sensitive shuttle vectors containing *gltA* gene were named pSFKTC1, pSFKTC2 and pSFKTC4, respectively.

(6) Introduction of mutant *gltA* promoter into chromosome:

[0083] pSFKTC1, pSFKTC2 and pSFKTC4 were each introduced into *Brevibacterium lactofermentum* FGR2 strain by electrical pulse method. The transformants were selected on CM2B medium plates containing 25 μ g/ml of kanamycin at 25°C. After introduction of each plasmid, each obtained strain was cultured in CM2B liquid medium, spread on CM2B plates containing 25 μ g/ml of kanamycin, after the dilution to a concentration of 10^3 to 10^5 cfu per plate and cultured at 34°C. The strain having the temperature-sensitive plasmid became sensitive to kanamycin because the replication of the plasmid was inhibited at this temperature and, therefore, it could not form colonies. On the other hand, the strain having plasmid DNA integrated into the chromosome could be selected because it formed the colonies. Colonies thus obtained were taken and separated into respective colonies. Chromosomal DNA was extracted from the strain. PCR was conducted by using the chromosomal DNA as the template and primers of sequence shown in Seq ID No. 8 and Seq ID No. 13. About 3 kb of amplified fragments were confirmed. It was thus proved that in this strain, mutant *gltA* gene derived from the temperature-sensitive plasmid was integrated near *gltA* gene in the host chromosome by homologous recombination. Strains derived from pSFKTC1, 2 and 4 were named BLCS11, BLCS12 and BLCS14, respectively.

(7) Preparation of substituted *gltA* promoters:

[0084] First, kanamycin-sensitive strains were obtained from the strains BLCS11, BLCS 12 and BLCS14 having mutant *gltA* gene integrated therein by homologous recombination. The strains having plasmid integrated therein were diluted and spread on CMM2B plates and then cultured at 34°C. After the formation of colonies, the replicas of the plates were made using CM2B plates containing 25 μ g/ml of kanamycin and were cultured at 34°C. Thus, kanamycin-sensitized strains were obtained.

[0085] The chromosome was extracted from the kanamycin sensitive strain, and PCR was conducted with primers

having the sequence shown in Seq ID No. 7 and Seq ID No.8 to prepare *gltA* gene fragments. The amplified fragments thus obtained were purified with SUPRECO2 (Takara shuzo Co., Ltd.) and then subjected to the sequencing reaction using a primer of Seq ID No. 13 to determine the sequence in the promoter region thereof. As a result, the strain having the same promoter sequence as that of pKF19CS1 in Table 7 was named GB01, the strain having the same promoter sequence as that of pKF19CS2 was named GB02 and the strain having the same promoter sequence as that of pKF19CS4 was named GB03. It was indicated that in these strains, the *gltA* gene of wild type originally located on the chromosome was excised together with the vector plasmid while the mutant *gltA* gene introduced by the plasmid was remained on the chromosome when the plasmid and duplicated *gltA* gene were excised from the chromosome.

(8) Determination of activity of citrate synthase of mutant *gltA* promoter strains:

[0086] The activities of the citrate synthase were determined by treating FGR2, GB01, GB02, GB03 and FGR2/pSFKC strains obtained in step (7) in the same manner as that of step (4). The results are shown in Table 10. It was confirmed that the citrate synthase activity of the substituted *gltA* promoter strain was higher than that of the parent strains thereof.

Table 10

Strain	dABS/min/mg	Relative activity
FGR2	7.9	1.0
GB01	9.5	1.2
GB02	15.0	1.9
GB03	31.6	4.0
FGR2/pSFKC	61.6	7.8

(9) Results of culture of substituted *gltA* promoter strains:

[0087] Each of the strains obtained in above-described step (7) was inoculated into a seed culture medium having a composition shown in Table 11, and the culture was shaken at 31.5°C for 24 hours. 300 ml of a main culture medium having a composition shown in Table 11 was placed into 500 ml glass jar fermenters and then sterilized by heating and was inoculated by 40 ml of the seeds cultured as described above. The culture was started at a culture temperature of 31.5°C while the stirring rate and the aeration rate were controlled at 800 to 1300 rpm and 1/2 to 1/1 vvm, respectively. The pH of the culture liquid was kept at 7.5 with gaseous ammonia. The temperature was shifted to 37°C 8 hours after the initiation of the culture. The culture was terminated when glucose had been completely consumed in 20 to 40 hours, and the quantity of L-glutamic acid formed and accumulated in the culture liquid was determined.

[0088] As a result, the larger improvement in the yield of L-glutamic acid was confirmed when each of the strains GB02 and GB03 rather than GB01 and FGR2/pSFKC was used as shown in Table 12. From these facts, it was found that good results were obtained by introducing the mutation into the *gltA* promoter to increase the CS activity to 2 to 4 times for the improvement in the yield of glutamic acid produced by those strains.

Table 11

Ingredient	Concentration	
	Seed culture	Main culture
Glucose	50 g/l	150 g/l
KH ₂ PO ₄	1 g/l	2 g/l
MgSO ₄ ·7H ₂ O	0.4 g/l	1.5 g/l
FeSO ₄ ·7H ₂ O	10 mg/l	15 mg/l
MnSO ₄ ·4H ₂ O	10 mg/l	15 mg/l
Soybean protein hydrolyzate	20 ml/l	50 ml
Biotin	0.5 mg/l	2 mg/l

Table 11 (continued)

Ingredient	Concentration	
	Seed culture	Main culture
Thiamine hydrochloride	2 mg/l	3 mg/l

Table 12

Strain	L-glutamic acid (g/l)
FGR2	8.9
GB01	9.1
GB02	9.4
GB03	9.4
FGR2/pSFKC	9.1

Example 4 Introduction of mutation into ICDH gene promoter region of coryneform glutamate-producing bacterium:

[0089] In this Example, strains having enhanced promoters for genes which codes glutamate dehydrogenase, citrate synthase and isocitrate dehydrogenase were produced.

(1) Cloning of *icd* gene:

[0090] The DNA sequence of *icd* gene of coryneform bacterium, which codes citrate synthase, has already been elucidated [J. Bacteriol. 177, 774-782 (1995)]. On the bases of this sequence, primers shown in Seq ID No. 14 and Seq ID No.15 were synthesized. PCR was conducted by using chromosomal DNA of *Brevibacterium lactofermentum* ATCC13869 as the template to amplify about 3 Kbp of DNA fragment containing *icd* gene and promoter thereof. The amplified fragment thus obtained was completely digested with *Eco*RI, and mixed with that obtained by complete digestion of pHSG399 (Takara Shuzo Co., Ltd.) with *Eco*RI to conduct the ligation. After the completion of the ligation, the transformation was conducted using competent cells of *E. coli* JM109. The cells were spread on the L-medium plates containing 10 µg/ml of IPTG, 40 µg/ml of X-Gal and 40 µg/ml of chloramphenicol. After overnight incubation, white colonies were taken and the transformants was obtained after single colony isolation.

[0091] The plasmid carrying *icd* gene was named pHSG399*icd*.

(2) Introduction of mutations into *icd* promoter:

[0092] The accurate location of the promoter of *icd* gene has not yet been determined. The possibility of increasing mRNA transcription level of *icd* gene was investigated by artificially modifying the upstream sequence of the gene which codes ICDH into a promoter-like sequence. Specifically, mutations were introduced into the -10 like region existing in the DNA sequence about 190 bp upstream (the first promoter) and about 70 bp (the second promoter) upstream from the first ATG of ICDH protein.

[0093] Mutan-Super Express Km (Takara Shuzo Co., Ltd.) was used for the introduction of mutation into an upstream region of *icd* gene. The method is specifically described below. pHSG399*icd* was completely digested with *Pst*I to obtain *Pst*I fragment containing the promoter of *icd* gene. The fragments were ligated with the fragment obtained by complete digestion of pKF18*icd* (Takara Shuzo Co., Ltd.) with *Pst*I. After the completion of the ligation, the transformation was conducted with competent cells of *E. coli* JM109 (Takara Shuzo Co., Ltd.). The cells was spread on the L-medium containing 10 µg/ml of IPTG, 40 µg/ml of X-Gal and 25 µg/ml of kanamycin. After overnight incubation, white colonies were taken and transformants were obtained after single colony isolation. From the transformants, plasmids were prepared, and the plasmid containing the promoter of *icd* gene was named pKF18*icd*.

[0094] PCR was conducted by using pKF18*icd* as the template and 5'-phosphorylated synthetic DNA shown in Seq ID No. 16, Seq ID No.17, Seq ID No.18, Seq ID No.19, Seq ID No.20 and Seq ID No.21 and the selection primer. These PCR products were used for transforming competent cells of *E. coli* JM109. The cells were spread on the L-medium

plates containing 25 μ g/ml of kanamycin. After overnight incubation, formed colonies were taken and the transformants were obtained after single colony isolation. From the transformants, plasmid DNA was prepared, and the sequence of *icd* promoter region was determined using synthetic DNA shown in Seq ID No. 22 by Sanger's method [J. Mol. Biol., 143, 161 (1980)]. Specifically, the DNA sequence was determined with Dye Terminator Sequencing Kit (Applied Biosystems), and analyzed with Genetic Analyzer ABI310 (Applied Biosystems). Those obtained by replacing *icd* promoter region with a sequence shown in Table 7 were named pKF18ICD1, pKF18ICD2, pKF18ICD3, pKF18ICD4, pKF18ICD5 and pKF18ICD6m respectively. Among them, pKF18ICD2 was completely digested with PstI to obtain PstI fragment containing the promoter of *icd* gene. The fragment was ligated with the fragment obtained by complete PstI digestion of pKF18kM (Takara Shuzo Co., Ltd.). After the completion of the ligation, the transformation was conducted with competent cells of *E. coli* JM109 (Takara Shuzo Co., Ltd.). The cells were spread on the L-medium plates containing 10 μ g/ml of IPTG, 40 μ g/ml of X-Gal and 25 μ g/ml of kanamycin. After overnight incubation, white colonies were taken and the transformed strains were obtained after single colony isolation. From the transformed strains, plasmids were prepared, and the plasmid containing the promoter of *icd* gene was named pKF18ICDM2. PCR was conducted using pKF18ICDM2 as the template and 5'-phosphorilated synthetic DNA shown in Seq ID No. 20 and Seq ID No.21 and the selection primer. The transformation of competent cells of *E. coli* JM109 was conducted with the PCR product. The cells were spread on the L-medium plates containing 25 μ g/ml of kanamycin. After overnight incubation, colonies thus formed were taken and transformants were obtained after single colony isolation. From the transformants, plasmids DNA were prepared, and the sequence of *icd* promoter region was determined using synthetic DNA shown in Seq ID No. 22. Those obtained by replacing *icd* promoter region with the sequence shown in Table 13 were named pKF18ICD25 and pKF18ICD26, respectively.

Table 13

Plasmid	1st Promoter		2nd Promoter	
	-35	-10	-35	-10
pKF18ICD	GCGACT	GAAAGT	TTTCCA	CACCAT
pKF18ICD01	GCGACT	TATAAT	TTTCCA	CACCAT
pKF18ICD02	TTGACA	TATAAT	TTTCCA	CACCAT
pKF18ICD03	TTGACT	TAAAGT	TTTCCA	CACCAT
pKF18ICD04	GCGACT	GAAAGT	TTTCCA	TATAAT
pKF18ICD05	GCGACT	GAAAGT	TTGCCA	TATAAT
pKF18ICD06	GCGACT	GAAAGT	TTGACA	TATAAT
pKF18ICD25	TTGACA	TATAAT	TTGCCA	TATAAT
pKF18ICD26	TTGACA	TATAAT	TTGACA	TATAAT

(3) Plasmid construction for determination of promoter activity:

[0095] For easily determining the promoter activity, a possible method is the indirect determination of the promoter activity using a reporter gene. Desirable properties required of the reporter gene are that the activity can be easily determined, that even when an amino acid is added to an N-terminal, the activity is not seriously lowered, that the background reaction does not occur and that it has a restriction enzyme cleavage site suitable for the gene manipulation. Because β galactosidase (LacZ) of *E. coli* is widely used as a reporter gene and bacteria of the genus *Corynebacterium* do not have lactose assimilability [J. Gen. Appl. Microbiol., 18, 399-416 (1972)], LacZ was determined to be the optimum reporter gene. Then, plasmid pNEOL carrying LacZ as the reporter gene was constructed (see Fig.3). The process for the construction is described in detail below. PCR was conducted by using a chromosomal DNA obtained from *E. coli* ME8459 (ME8459 was deposited with National Institute of Genetics (Japan)) as the template with synthetic DNA shown in Seq ID No. 23 and Seq ID No.24 as the primer. The PCR product was completely digested with SmaI and BamHI and then ligated with fragments obtained by digesting pKF3 (Takara Shuzo Co., Ltd.) with HindIII and blunt-ended. After the completion of the ligation, the transformation was conducted with competent cells of *E. coli* JM109 (Takara Shuzo Co., Ltd.). The cells were spread on the L-medium plates containing 25 μ g/ml of kanamycin. After overnight incubation, colonies thus formed were taken and separated into respective colonies to obtain the transformed strain. The plasmid obtained from the transformed strain was named pKF3nptII. Then, this plasmid was digested with Sall. On the other hand, pSAK4 described in Example 1(1) was completely digested with SmaI and Sall and blunt-

ended. These fragments were ligated together to construct a shuttle vector pNEO which can replicate in a coryneform bacterium. This plasmid was capable of imparting a resistance to chloramphenicol and resistance to kanamycin to the hosts. Further, pNEO was completely digested with *Sma*I and *Sse*8387I. The resultant fragments were ligated to those obtained by complete digestion of pMC1871 (Farmacia Biotech.) with *Pst*I and *Sma*I. Thus, shuttle vector pNEOL which can be replicated in a coryneform bacterium and having LacZ lacking 8 amino acid on N-terminal as the reporter gene was constructed (see Fig.3).

(4) Determination of activity of mutant *icd* promoter:

[0096] Plasmids having mutant *icd* promoter constructed in above-described step (2), i. e. pKF18ICD1, pKF18ICD2, pKF18ICD3, pKF18ICD4, pKF18ICD5, pKF18ICD6, pKF18ICD25, pKF18ICD26 and pKF18ICD, were completely digested with *Sac*II and *Pst*I and then blunt-ended. They were ligated with fragment obtained by digesting pNEOL with *Sma*I. After the completion of the ligation, the transformation was conducted with competent cells of *E. coli* JM109. The cells were spread on the L-medium plates containing IPTG, X-Gal and 40 μ g/ml of chloramphenicol. After overnight incubation, blue colonies were taken and the transformed strains were obtained after single colony isolation.

[0097] From the transformed strains, plasmids were prepared. Plasmids having a structure capable of producing a fused protein of ICDH and LacZ were named pNEOICD1, pNEOICD2, pNEOICD3, pNEOICD4, pNEOICD5, pNEOICD6, pNEOICD25, pNEOICD26 and pNEOLICD, respectively. Each of These plasmids or pNEOL was introduced into *Brevibacterium lactofermentum* ATCC13869 by electrical pulse method. The transformants were selected by using CM2B medium plates (comprising 10 g/l of bactotryptone, 10 g/l of bacto yeast extract, 5 g/l of NaCl, 10 μ g/l of biotin and 15 g/l of agar and having pH 7.0) containing 25 μ g/ml of kanamycin and 40 μ g/ml of X-Gal at 31 °C for two days. After the completion of the introduction, colonies thus formed were taken and isolated as single colonies. The transformants containing pNEOICD1, pNEOICD2, pNEOICD3, pNEOICD4, pNEOICD5, pNEOICD6, pNEOICD25, pNEOICD26 and pNEOLICD were named BLAC1, BLAC2, BLAC3, BLAC4, BLAC5, BLAC6, BLAC25, BLAC26, BLAC and BNEOL, respectively. All the transformants other than BNEOL formed blue colonies. Crude enzyme solutions were prepared from the transformants in the same manner as that of step (4) in Example 3 except that "Z-Buffer" (comprising 10 mM of KCl, 1 mM of $MgSO_4$, 270 μ g/100 mM of 2-ME and NaPi and having pH 7.5) was used as a washing and suspension buffer. The activity of LacZ was determined as follows: Z-Buffer was mixed with the crude enzyme solution, ONPG in Z-Buffer having the final concentration of 0.8 mg/ml was added to the resultant mixture, and the increase in the absorbance at 420 nm at 30°C was determined with Hitachi spectrophotometer U-3210 as the activity of LacZ. The protein concentration in the crude enzyme solution was determined by Protein Assay (BIO-RAD). Bovine serum albumin was used as the standard protein. The results are shown in Table 14. It was confirmed that the LacZ activity of the strain having a mutation in *icd* promoter and expressing ICDH-LacZ fused protein was higher than that expressing the wild type ICDH-LacZ fused protein.

Table 14

Strain	dABS/min/mg	Relative activity
BNEOL	Not detected	0.0
BNEOL1	42	1.0
BNEOL1-1	84	2.0
BNEOL1-2	168	4.0
BNEOL1-3	80	1.9
BNEOL1-4	126	3.0
BNEOL1-5	139	3.3
BNEOL1-6	84	2.0
BNEOL1-25	168	4.0
BNEOL1-26	170	4.0

(5) Introduction of mutant *icd* gene into temperature-sensitive plasmid:

[0098] Plasmid vector pSFKT2 (Japanese Patent Application No. 11-81693) the replication of which in a coryneform bacterium was temperature-sensitive was used. pKF18ICD1, pKF18ICD2, pKF18ICD3, pKF18ICD4, pKF18ICD5,

pKF18ICD6, pKFICD25 and pKFICD26 were completely digested with PstI and the obtained fragments were used as the mutant *icd* promoter sequences. The fragments thus obtained were ligated with pSFKT2 completely digested with PstI. After the completion of the ligation, the transformation was conducted with competent cells of *E. coli* JM109 (Takara Shuzo Co., Ltd.). The cells were spread on the L-medium plates containing 10 μ g/ml of IPTG, 40 μ g/ml of X-Gal and 25 μ g/ml of kanamycin. After overnight incubation, white colonies were taken and transformed strains were obtained after single colony isolation. From the transformed strains, plasmids were prepared. Temperature-sensitive shuttle vectors containing *icd* promoter were named pSFKT11, pSFKT12, pSFKT13, pSFKT14, pSFKT15, pSFKT16, pSFKT125 and pSFKT126, respectively.

(6) Integration of mutant *icd* promoter into chromosome:

[0099] The plasmids constructed in above-described step (5) were each introduced into *Brevibacterium lactofermentum* GB02 strain by electrical pulse method. The transformants were selected with CM2B medium plates (comprising 10 g/l of bactotryptone, 10 g/l of bacto yeast extract, 5 g/l of NaCl, 10 μ g/l of biotin and 15 g/l of agar and having pH 7.0) containing 25 μ g/ml of kanamycin at 25°C. After the completion of the introduction, the obtained strains were cultured in CM2B liquid medium, spread on CM2B plates containing 25 μ g/ml of kanamycin after the dilution to a concentration of 10^3 to 10^5 cfu per plate and cultured at 34°C. The strain having the temperature-sensitive plasmid became sensitive to kanamycin because the replication of the plasmid was inhibited at this temperature and, therefore, it could not form colonies. On the other hand, the strain having plasmid DNA integrated into the chromosome could be selected because it could form colonies. Colonies thus obtained were taken and separated into isolated colonies. Chromosomal DNA was extracted from the strain and PCR was conducted by using the chromosomal DNA as the template with primers shown in Seq ID No. 13 and Seq ID No. 15. About 3 kb of amplified fragments were confirmed. It was thus proved that in this strain, mutant *icd* gene derived from the temperature-sensitive plasmid was integrated near *icd* gene in the host chromosome by homologous recombination.

(7) Preparation of strains having substituted *icd* promoter:

[0100] First, kanamycin-sensitive strain was obtained from the strains having mutant *icd* gene integrated therein by the homologous recombination as described in step (6). The strains having the plasmid integrated therein were diluted and spread on CM2B plates and then cultured at 34°C. After the formation of colonies, replicas were made on CM2B plates containing 25 μ g/ml of kanamycin, and they were incubated at 34°C. Thus, kanamycin-sensitive strains were obtained.

[0101] The chromosome was extracted from the kanamycin resistant strain, and PCR was conducted using primers shown in Seq ID No. 14 and Seq ID No. 15 to prepare *icd* gene fragments. The amplified fragments thus obtained were purified with SUPRECO2 (Takara Shuzo Co., Ltd.) and then subjected to the sequencing reaction using a primer shown in Seq ID No. 22 to determine the sequence of the promoter region thereof. As a result strains having *icd* promoter sequences derived from pSFKT11, pSFKT12, pSFKT13, pSFKT14, pSFKT15, pSFKT16, pSFKT125 and pSFKT126 were named GC01, GC02, GC03, GC04, GC05, GC06, GC25 and GC26, respectively. In these strains, when the plasmid and duplicate *icd* gene were excised from the chromosome, the *icd* gene of wild type originally located on the chromosome was excised together with the vector plasmid, while the mutant *icd* gene introduced by the plasmid remained on the chromosome.

(8) Determination of isocitrate-dehydrogenase activity of the mutant strains having mutant *icd* promoter:

[0102] ICDH crude enzyme solution was prepared by using each of the 8 strains obtained in above-described step (7) and GB02 strain in the same manner as that of step (7) in Example 3. The ICDH activities were determined as follows: The crude enzyme solution was added to a reaction solution containing 35 mM of TrisHCl (pH 7.5), 1.5 mM of $MnSO_4$, 0.1 mM of NADP and 1.3 mM of isocitric acid, and the increase in the absorbance at 340 nm at 30°C was determined with Hitachi spectrophotometer U-3210 as the activity of ICDH. The protein concentration in the crude enzyme solution was determined by Protein Assay (BIO-RAD). Bovine serum albumin was used as the standard protein. The results are shown in Table 15. It was confirmed that the isocitrate dehydrogenase activity of substituted *icd* promoter strains was higher than that of the parent strain.

Table 15

Strain	dABS/min/mg	Relative activity
GB02	3.9	1.0

Table 15 (continued)

Strain	dABS/min/mg	Relative activity
GC01	8.2	2.1
GC02	19.1	4.9
GC03	7.0	1.8
GC04	12.5	3.2
GC05	19.1	4.9
GC06	10.5	2.7
GC25	30.4	7.8
GC26	24.2	6.2

(9) Results of culturing the strains containing substituted *icd* promoter:

[0103] Each of the 8 strains obtained in above-described step (7) was cultured in the same manner as that in step (9) in Example 3. As a result, the improvement in the yield of L-glutamic acid was confirmed when any one of the strains GC02, GC04, GC05, GC25 and GC26 was used as shown in Table 16. It was found that good results were obtained by introducing the mutation into *icd* promoter to increase the ICDH activity to at least 3 times.

Table 16

Strain	L-glutamic acid (g/dl)
GB02	9.2
GC01	9.0
GC02	9.5
GC03	9.1
GC04	9.4
GC05	9.6
GC06	9.2
GC25	9.9
GC26	9.8

Example 5 Introduction of mutation into PDH gene promoter region of coryneform glutamate-producing bacterium:

(1) Cloning of *pdhA* gene from coryneform bacteria

[0104] Primers shown in Seq ID No.25 and Seq ID No.26 were synthesized by selecting regions having a high homology among E1 subunits of pyruvate dehydrogenase (PDH) of *Escherichia coli*, *Pseudomonas aeruginosa* and *Mycobacterium tuberculosis*. PCR was conducted by using chromosome of *Brevibacterium lactofermentum* ATCC13869, prepared with a bacterial genomic DNA purification kit (Advanced Genetic Technologies Corp.), as the template under standard reaction conditions described on page 8 of PCR Technology (edited by H. Erlich and published by Stockton Press, 1989). The reaction solution was subjected to the electrophoresis in an agarose gel to find that about 1.3 kilobases of DNA fragment was amplified. The sequence of both end of the obtained DNA was determined with synthetic DNA shown in Seq ID No. 25 and Seq ID No.26. The sequence was determined by Sanger's method [J. Mol. Biol., 143, 161 (1980)] with DNA Sequencing Kit (Applied Biosystems Co.). The determined sequence was deduced to amino acids, and compared with E1 subunits of pyruvate dehydrogenase derived from each of *Escherichia coli*, *Pseudomonas aeruginosa* and *Mycobacterium tuberculosis* to find a high homology among them. Consequently, it was determined that the DNA fragment amplified by PCR was a part of *pdhA* gene which codes E1 subunit of pyruvate

dehydrogenase of *Brevibacterium lactofermentum* ATCC13869. The cloning of the upstream and downstream region of the gene was conducted. The cloning method was as follows: A chromosome of *Brevibacterium lactofermentum* ATCC13869 was digested with restriction enzymes EcoRI, BamHI, Hind III, Pst I, Sal I and Xba I (Takara Shuzo Co., Ltd.) to obtain DNA fragments. LA PCR in vitro cloning Kit (Takara Shuzo Co., Ltd.) was used for the cloning, using the sequences shown in Seq ID No. 27 and Seq ID No.28 in the Sequence Listing as primers for cloning the upstream region, and sequences shown in Seq ID No. 29 and Seq ID No.30 as primers for cloning the downstream region. After PCR using the kit, DNA fragments of about 0.5, 2.5, 3.0, 1.5 and 1.8 kilobases were amplified for the upstream region from the fragments obtained by digestion with EcoRI, Hind III, Pst I, Sal I and Xba I, respectively; and DNA fragments of about 1.5, 3.5 and 1.0 kilobase were amplified for the downstream region from the fragments obtained by digestion with BamHI, Hind III and Pst I, respectively. The sequences of these DNA fragments were determined in the same manner as that described above. It was found that the amplified DNA fragments further contained an open reading frame of about 920 amino acids and also that a region supposed to be a promoter region was present in the upstream region. Because the deduced amino acid sequence from the DNA sequence of the open reading frame is highly homologous to known E1 subunit of pyruvate dehydrogenase such as that of *E. coli*, it was apparent that the open reading frame was the *pdhA* gene which codes E1 subunit of pyruvate dehydrogenase of *Brevibacterium lactofermentum* ATCC13869. The DNA sequence of the open reading frame was shown in Seq ID No. 31 in the Sequence Listing. In Seq ID No. 31 in the Sequence Listing, deduced amino acid sequence from the DNA sequence is also shown. Since methionine residue at N-terminal of the protein is derived from ATG which is an initiation codon, it usually does not concern the essential function of protein, and it is well known that the methionine residue is removed by the effect of peptidase after the translation. Therefore, in the above-described protein, it is possible that methionine residue at the N-terminal has been removed. However, the GTG sequence is present in 6 bases upstream of ATG shown in Seq ID No. 31 in the Sequence Listing, and it is also possible that amino acids is translated from this point. Pyruvate dehydrogenase of other microorganisms such as *E. coli* are composed of three subunits of E1, E2 and E3, and genes which encode them constitute an operon in many cases. However, there was no open reading frame considered to be E2 and E3 subunit of pyruvate dehydrogenase in about 3 kilobases downstream of *pdhA* gene. Instead, it was shown that a sequence supposed to be a terminator was present in the downstream of the open reading frame. From these facts, it was supposed that E2 and E3 subunits of pyruvate dehydrogenase of *Brevibacterium lactofermentum* ATCC13869 were present in another region on the chromosome.

(2) Construction of a plasmid for amplifying *pdhA*:

[0105] It was already apparent that a strain obtained by introducing a gene which codes three subunits constituting PDH of *E. coli* into *Brevibacterium lactofermentum* ATCC13869 gives an improved glutamic acid yield (JP No.10-360619). However, in PDH of *Brevibacterium lactofermentum* ATCC13869, only *pdhA* gene which codes E1 subunit had been cloned, and no examination had not been made to know whether the amplification of the gene alone is effective in improving the yield of glutamic acid. Under these circumstances, examination was made to know whether the amplification of *pdhA* gene alone is effective in improving the yield of glutamic acid or not.

[0106] Primers shown in Seq ID No. 33 and Seq ID No.34 were synthesized on the basis of the previously cloned DNA sequences. PCR was conducted by using chromosome of *Brevibacterium lactofermentum* ATCC13869, prepared with a Bacterial Genomic DNA Purification kit (Advanced Genetic Technologies Corp.), as the template under standard reaction conditions described on page 8 of PCR Technology (edited by H. Erlich and published by Stockton Press, 1989) to amplify *pdhA* gene. Among the primers thus synthesized, Seq ID No. 33 corresponded to a sequence of base No. 1397 to No.1416 in *pdhA* gene described in Seq ID No. 32 in the Sequence Listing. Seq ID No. 34 was the complementary strand of the DNA sequence corresponding to the sequence of base No. 5355 to No.5374 in Seq ID No. 32 in the Sequence Listing, which was represented from the 5' side.

[0107] PCR product thus obtained was purified by an ordinary method and reacted with restriction enzyme Sal I and EcoT22I. The fragment was ligated with pSFK(Patent Application No.11-69896), cleaved with restriction enzymes Sal I and Pst I, with a ligation kit (Takara Shuzo Co., Ltd.). After the transformation with competent cells (Takara Shuzo Co., Ltd.) of *E. coli* JM109, the cells were spread to the L-medium medium plates(comprising 10 g/l of bactotryptone, 5 g/l of bacto yeast extract, 5 g/l of NaCl and 15 g/l of agar and having pH 7.2) containing 10 μ g/ml of IPTG (isopropyl- β -D-thiogalactopyranoside), 40 μ g/ml of X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) and 25 μ g/ml of kanamycin. After overnight incubation, white colonies were taken and the transformed strains were obtained after single colony isolation.

[0108] From the transformed strains, plasmids were prepared by alkali method (*Seibutsu Kogaku Jikken-sho* edited by Nippon *Seibutsu Kogaku kai* and published by *Baifukan*, p. 105, 1992). Restriction enzyme maps of DNA fragments inserted into the vectors were prepared and compared with the restriction enzyme map of *pdhA* gene reported in sequence No. 32 of the Sequence Listing. A plasmid containing DNA fragments inserted therein having the same restriction enzyme map as that of *pdhA* gene was named pSFKBPDHA.

(3) Introduction of pASFKBDHA into *Brevibacterium lactofermentum* ATCC13869 and GC25 and evaluation of the fermentation experiments:

[0109] *Brevibacterium lactofermentum* ATCC13869 and GC25 were transformed with plasmid pSFKBDHA by electrical pulse method (J. P. KOKAI No.2-207791) to obtain the transformed strains. The culture for producing L-glutamic acid was conducted with transformed strain ATCC13869/pSFKBDHA and GC25/pSFKBDHA obtained by introducing plasmid pSFKBDHA into *Brevibacterium lactofermentum* ATCC13869 and GC25 as follows: Cells of ATCC13869/pSFKBDHA and GC25/pSFKBDHA obtained by the culture on CM2B medium plates containing 25 μ g/ml of kanamycin were inoculated into a medium (comprising 1 liter of pure water containing 80 g of glucose, 1 g of KH_2PO_4 , 0.4 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 30 g of $(\text{NH}_4)_2\text{SO}_4$, 0.01 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01 g of $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$, 15 ml of soybean protein hydrolyzate, 200 μ g of thiamine hydrochloride, 60 μ g of biotin, 25 mg of kanamycin and 50 g of CaCO_3 ; and having a pH adjusted to 8.0 with KOH). Then the culture was shaken at 31.5°C until sugar in the medium had been consumed. The obtained products were inoculated into the medium of the same composition as that described above (for GC25/pSFK6 and GC25/pSFKBDHA) or the medium eliminated Biotin from the composition as that described above (for ATCC13869/pSFK6 and ATCC13869/pSFKBDHA) in an amount of 5 %, and the shaking culture was conducted at 37°C until sugar in the medium had been consumed. As a control, strains obtained by transforming *Brevibacterium lactofermentum* ATCC13869 and GC25 with previously obtained plasmid pSFK6 capable of autonomously replicating in coryneform bacterium by electrical pulse method (J. P. KOKAI No. 2-207791), were cultured in the same manner as that described above. After the completion of the culture, the amount of L-glutamic acid accumulated in the culture medium was determined with Biotic Analyzer AS-210 (a product of Asahi Chemical Industry Co., Ltd.). The results are shown in Table 17.

Table 17

Strain	Yield of L-glutamic acid (g/dl)
ATCC13869/Psfk	3.6
ATCC13869/pSFKBDHA	3.8
GC25/pSFK 6	5.1
GC25/pSFKBDHA	5.3

[0110] From these results, it was apparent that even the amplification of *pdhA* gene alone is sufficiently effective in improving the yield of Glu in *Brevibacterium lactofermentum* ATCC13869 and GC25.

(4) Construction of plasmids for determination of the activity of mutated *pdhA* promoter:

[0111] To produce promoter mutant of pyruvate dehydrogenase (PDH), the determination of the previously cloned promoter region of *pdhA* gene of *Brevibacterium lactofermentum* ATCC13869 was conducted and also the determination of difference in the expression caused by the modification of the promoter region were conducted by determining the activity of β -galactosidase.

[0112] The promoter region of *pdhA* gene was presumed from the DNA sequence which had been already elucidated by cloning. As a result, it was supposed to be possible that base No. 2252 to No.2257 and No. 2279 to No. 2284 in Seq ID No. 32 in the Sequence Listing were -35 region and -10 region, respectively. Therefore, primers shown as Seq ID No. 35 and Seq ID No.36 in the Sequence Listing were synthesized, and DNA fragments containing promoter region of *pdhA* gene were amplified by PCR method by using chromosomal DNA of *Brevibacterium lactofermentum* ATCC13869 as a template. Among the synthesized primers, Seq ID No. 35 corresponded to the sequence ranging from base No. 2194 to base No. 2221 in Seq ID No. 32; but the base No. 2198 had been replaced with C, and the base No. 2200 and No.2202 had been replaced with G, and recognition sequence for restriction enzyme *Sma*I had been inserted. Seq ID No. 36 corresponded to the sequence ranging from base No. 2372 to base No. 2398 in Seq ID No. 32; but base No.2393 and No.2394 had been replaced with G, and the complementary strand of the DNA sequence having a recognition sequence of restriction enzyme *Sma*I inserted therein was represented from the 5'-end. PCR was conducted by using chromosome of *Brevibacterium lactofermentum* ATCC13869, prepared with Bacterial Genomic DNA Purification kit (Advanced Genetic Technologies Corp.), as the template under standard reaction conditions described on page 8 of PCR Technology (edited by H. Erlich and published by Stockton Press, 1989) to amplify the promoter region of *pdhA* gene. PCR product thus obtained was purified by an ordinary method and reacted with restriction enzyme *Sma*I. The

fragments were ligated with pNEOL lacking in promoter region of lacZ gene which could be replicate in a coryneform bacterium and which had been digested with restriction enzymes Sma I, (Example 4 (3) with a Ligation Kit (Takara Shuzo Co., Ltd.). After the transformation with competent cells (Takara Shuzo Co., Ltd.) of *E. coli* JM109, the cells were spread on the L-medium plates (comprising 10 g/l of bactotryptone, 5 g/l of bacto yeast extract, 5 g/l of NaCl and 15 g/l of agar and having pH 7.2) containing 40 μ g/ml of X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) and 25 μ g/ml of kanamycin. After overnight incubation, blue colonies were taken and the transformed strains were obtained after single colony isolation. From the transformants, plasmids were prepared by alkali method (*Seibutsu Kogaku Jikken-sho* edited by *Nippon Seibutsu Kogaku-kai* and published by *Baifukan*, p. 105, 1992). After sequencing DNA fragments inserted into the vector by an ordinary method, the plasmid containing the DNA fragment inserted therein was named pNEOLB-PDHAprol.

[0113] Further, primers indicated as Seq ID No. 37, Seq ID No.38 and Seq ID NO.39 in the Sequence Listing were synthesized for constructing plasmids wherein a region supposed to be the promoter site was changed to the consensus sequence of promoters of coryneform bacteria. By using each of the primers and a primer shown in Seq ID No. 36, DNA fragments wherein the promoter region of *pdhA* gene was changed to the consensus sequence were amplified by PCR method by using chromosomal DNA of *Brevibacterium lactofermentum* ATCC13869 as a template. Among the synthesized primers, Seq ID No. 37 corresponded to the sequence ranging from base No. 2244 to base No. 2273 in Seq ID No. 32; base No. 2255 had been replaced with C, and base No. 2257 had been replaced with A; thus only -35 region had been changed to the consensus sequence of the coryneform bacteria. Seq ID No. 38 corresponded to the sequence ranging from base No. 2249 to base No. 2288 in sequence No. 32; base No. 2279 and No.2281 had been replaced with T; thus only -10 region had been changed to the consensus sequence of the coryneform bacteria. Sequence No. 39 corresponded to the sequence ranging from base No. 2249 to base No. 2288 in Seq ID No. 32; base No. 2255 had been replaced with C, base No. 2257 had been replaced with A, and base No. 2279 and No.2281 had been replaced with T; thus both -35 region and -10 region had been changed to the consensus sequence of the coryneform bacteria. PCR was conducted by using chromosome of *Brevibacterium lactofermentum* ATCC13869, prepared with a Bacterial Genomic DNA Purification Kit (Advanced Genetic Technologies Corp.), as the template under standard reaction conditions described on page 8 of PCR Technology (edited by H. Erlich and published by Stockton Press, 1989) to amplify the promoter region of *pdhA* gene with these primers so that the promoter region was changed to the consensus sequence. PCR products thus obtained were purified by an ordinary method and reacted with restriction enzyme SmaI. The fragments were ligated with pNEOL lacking the promoter region of lacZ gene, which could replicate in a coryneform bacterium and which had been cleaved with restriction enzymes Sma I, with a Ligation Kit (Takara Shuzo Co., Ltd.). After the transformation with competent cells (Takara Shuzo Co., Ltd.) of *E. coli* JM109, the cells were spread on the L-medium plates (comprising 10 g/l of bactotryptone, 5 g/l of bacto yeast extract, 5 g/l of NaCl and 15 g/l of agar and having pH 7.2) containing 40 μ g/ml of X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) and 25 μ g/ml of kanamycin. After overnight incubation, blue colonies were taken and the transformed strains were obtained after single colony isolation. From the transformed strains, plasmids were prepared by the alkali method (*Seibutsu Kogaku Jikken-sho* edited by *Nippon Seibutsu Kogaku kai* and published by *Baifukan*, p. 105, 1992). After sequencing DNA fragments inserted into the vector by an ordinary method, the plasmid containing DNA fragments, wherein only the sequence in -35 region had been changed to the consensus sequence, inserted therein was named pNEOLBPDHApro35; the plasmid containing DNA fragments, wherein only the sequence in -10 region had been changed to the consensus sequence, was inserted therein was named pNEOLBPDHApro10; and the plasmid containing DNA fragments, wherein the sequences in both -35 region and -10 region had been changed to the consensus sequence, was inserted therein was named pNEOLBPDHApro3510.

(5) The evaluation of the mutated *pdhA* promoter activity:

[0114] *Brevibacterium lactofermentum* ATCC13869 was transformed with plasmids named pNEOLBPDHApro1, pNEOLBPDHApro10 and pNEOLBPDHApro3510 by electrical pulse method (J. P. KOKAI No. 2-207791) to obtain the transformed strains. β -galactosidase activity of the obtained transformants was determined by the method described in Example 4(4). After changing the sequence in the promoter region to the consensus sequence, β -galactosidase activities were as shown in Table 18, wherein the enzymatic activity of β -galactosidase having the promoter region of *pdhA* gene was given as 1.

Table 18

Strain	β -Galactosidase activity (relative value)
ATCC13869/pNEOLBPDHApro1	1

Table 18 (continued)

Strain	β -Galactosidase activity (relative value)
ATCC13869/pNEOLBPDHAp010	6
ATCC13869/pNEOLBPDHAp03510	7.5

[0115] These results indicate that the supposed promoter region was the promoter of *pdhA* gene and that the expression of *PdhA* can be changed (enhanced) by changing the sequence in this region into the consensus sequence. This fact indicates that the expression can be changed, without using plasmid, by changing the promoter region of *pdhA* gene.

(6) Construction of plasmid for preparation of promoter varied strain:

[0116] Since it had been proved that the expression of *pdhA* can be changed by introducing mutations into the promoter, plasmids for preparing a *pdhA* promoter modified strains were constructed. Three constructs of the plasmids for the promoter modified strains were constructed. They were plasmids wherein -35 region, -10 region and both of them were changed to the consensus sequence, respectively.

[0117] Primers shown in Seq ID No. 40 and Seq ID No.41 were newly synthesized on the basis of the DNA sequence which had already been cloned. Among synthesized primers, Seq ID No. 40 was the complementary strand of the DNA sequence corresponding to a sequence ranging from base No. 2491 to base No. 2521 in Seq ID No. 32, which was represented from the 5'-end, and to which a sequence comprising three A's followed by four T's at the 5' terminal. Seq ID No. 33 was the complementary strand of the DNA sequence corresponding to the sequence ranging from base No. 5020 to base No. 5039 of *pdhA* gene in Seq ID No. 32, which was represented from the 5'-end. PCR was conducted by using Seq ID No. 33 and Seq ID No.40 as the primers and chromosome of *Brevibacterium lactofermentum* ATCC13869, prepared with Bacterial Genomic DNA Purification Kit (Advanced Genetic Technologies Corp.), as a template under standard reaction conditions described on page 8 of PCR Technology (edited by H. Erlich and published by Stockton Press. 1989). Further, PCR was conducted by using Seq ID No. 39 and Seq ID No. 41 and chromosome of *Brevibacterium lactofermentum* ATCC13869 as a template. The PCR products thus obtained were purified by an ordinary method. PCR was conducted by using PCR products obtained by using Seq ID No. 33 and No.40, PCR products obtained by using Seq ID No. 39 and Seq ID No.41 and Seq ID No. 33 and 41 as the primers. The PCR condition was as follows: The concentration of these four DNA would be 10 μ M in the reaction cocktail and La taq (Takara Shuzo Co., Ltd.) was used without template. PCR products were purified by an ordinary method, and reacted with restriction enzyme Sal I and Xho I. The fragments thus obtained were ligated with fragments obtained by digesting temperature-sensitive plasmid pSFKT2 with SalI, which can replicate in a coryneform bacterium, by using Ligation Kit (Takara Shuzo Co., Ltd.). After the transformation with competent cells (Takara Shuzo Co., Ltd.) of *E. coli* JM109, the cells was spread on the L-medium plates (comprising 10 g/l of bactotryptone, 5 g/l of bacto yeast extract, 5 g/l of NaCl and 15 g/l of agar and having pH 7.2) containing 10 μ g/ml of IPTG (isopropyl- β -D-thiogalactopyranoside), 40 μ g/ml of X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) and 25 μ g/ml of kanamycin. After overnight incubation, white colonies were taken and transformants were obtained after single colony isolation. From the transformants, plasmids were prepared by The alkali method (*Seibutsu Kogaku Jikken-sho* edited by *Nippon Seibutsu Kogaku-kai* and published by *Baifukan*, p. 105, 1992). After sequencing DNA fragments inserted into the vector, the base sequence was compared with that of *pdhA* gene reported in sequence No. 32. The plasmid containing DNA fragments, wherein only the sequences in -35 region and -10 region of the promoter were changed to the consensus sequence of the coryneform bacteria, inserted therein was named pSFKTPDHAp03510.

[0118] A plasmid wherein -35 region of the promoter of *pdhA* gene had been changed to the consensus sequence of coryneform bacteria, and also plasmid wherein -10 region of the promoter of *pdhA* gene had been changed to the consensus sequence of coryneform bacteria were constructed in the same manner as that described above except that Seq ID No. 39 in the Sequence Listing was replaced with Seq ID No. 37 and 38, respectively. These plasmids were named pSFKTPDHAp035 and pSFKTPDHAp010, respectively.

(7) Preparation of promoter modified strains:

[0119] Strains having modified *pdhA* gene promoter were prepared by the homologous recombination by using the plasmid for preparing promoter varied strain constructed in the above-described step (6).

[0120] First, GC25 was transformed with plasmid pSFKTPDHAp03510 for preparing promoter modified strain by electrical pulse method (refer to J. P. KOKAI No. 2-207791). The cells were spread on CM2B medium plates (comprising

10 g/l of polypeptone, 10 g/l of bacto yeast extract, 5 g/l of NaCl, 10 μ g/ml of biotin and 15 g/l of agar, and having pH 7.2) and cultured at 25°C to obtain transformed strains. These transformants were cultured in CM2B liquid medium in a test tube overnight and then spread on CM2B medium plates containing 25 μ g/ml of kanamycin and cultured at 34°C to obtain a strain caused by once-recombination which contains plasmid pSFKTPDHpro3510 on its chromosome inserted by the homologous recombination. After single colony isolation, this strain was cultured in CM2B liquid medium in a test tube overnight. After the suitable dilution, it was spread on CM2B medium plates and cultured at 31.5°C. After the colonies began to appear, the replicas were made on CM2B medium plates containing 25 μ g/ml of kanamycin to obtain kanamycin-sensitive strains. Since two kinds of the strains, i.e. a strain having the sequence of wild type strain for the promoter region of pdhA gene and another strain having the mutation introduced therein, could be occurred, this region was sequenced. Thus, a promoter modified strain, wherein the mutation had been introduced into the promoter region of pdhA gene, was obtained. In this strain, -35 region and -10 region of promoter of pdhA gene had been changed to the consensus sequence of coryneform bacteria. This strain was named GD3510.

[0121] Strains wherein -35 region or -10 region of promoter of pdhA gene had been changed to the consensus sequence of coryneform bacteria were obtained in the same manner as that described above except that above described plasmid pSFKTPDHpro3510 for producing the promoter modified strain was replaced with plasmid pSFKTPDHpro35 and pSFKTPDHpro10 for producing promoter modified strains and they were named GD35 and GD10, respectively.

(8) Evaluation of the results of flask culture of pdhA gene promoter modified strains:

[0122] The flask culture for producing L-glutamic acid was conducted with three kinds of pdhA gene promoter modified strains obtained as described above. Each of the cells of the promoter modified strains GD3510, GD35, GD10 and GC25 obtained by the culture on CM2B medium plates was inoculated into a medium (comprising 1 liter of pure water containing 30 g of glucose, 1 g of KH_2PO_4 , 0.4 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 30 g of $(\text{NH}_4)_2\text{SO}_4$, 0.01 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01 g of $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$, 15 ml of soybean hydrolyzate, 200 μ g of thiamine hydrochloride, 60 μ g of biotin and 50 g of CaCO_3 ; and having a pH adjusted to 8.0 with KOH). Then the culture was shaken at 31.5°C until the sugar in the medium had been consumed. The obtained products were inoculated into the medium of the same composition as that described above in an amount of 5%, and the shaking culture was conducted at 37°C until sugar in the medium had been consumed. After the completion of the culture, the amount of L-glutamic acid accumulated in the culture liquid was determined with Biotic Analyzer AS-210 (a product of Asahi Chemical Industry Co., Ltd.). The results are shown in Table 19.

Table 19

Strain	L-glutamic acid (g/dl)
GC25	1.9
GD35	2.0
GD10	2.0
GD3510	2.1

[0123] It was apparent from the results that the obtained promoter modified strains provided improved Glu yields.

Example 6 Introduction of mutation into promoter region of arginosuccinate synthase gene:

1) Determination of DNA sequence in the upstream of argG gene:

[0124] In order to amplify argG gene of *Brevibacterium flavum* by PCR, the DNA sequences in the upstream and downstream regions of the ORF were determined. The determination of the DNA sequences was conducted by synthesizing a primer based on the known DNA sequence (Gen Bank accession AF030520) of ORF of argG gene of *Corynebacterium glutamicum* and using in vitro LA PCR cloning kit (Takara shuzo Co., Ltd.) in accordance with the instruction manual included in the kit. As primers, they were specifically used oligonucleotide (primers 1 and 2) having the DNA sequences set out as Seq ID No. 42 and Seq ID No.43 for the upstream region, and oligonucleotide (primers 3 and 4) having the DNA sequences set out as Seq ID No.44 and Seq ID No.45 for the downstream region. The DNA sequences in the upstream and downstream region of argG were determined by completely digesting chromosome DNA of 2247 strain (ATCC14067), i.e., wild type strain of *Brevibacterium flavum*, with a restriction enzyme EcoRI, conducting first PCR with the primer 2 or 3 (having sequence No.43 or 44), and conducting second PCR with the primer 1 or 2 (having

sequence No. 42 or 45).

2) Prediction of promoter region:

- 5 [0125] A promoter-like sequence in the upstream of ORF of argG gene was search for the above-described sequences with a commercially available software (GENETYX). The mutation was introduced into a region of the highest score (about 120 bp upstream of the first ATG). Then, the promoter activity was measured.

3) Introduction of mutations into promoter sequence, and determination of activity of mutant promoters:

10

[0126] Mutation-introducing primers 9, 10, 11, 12 or 13 and 7 (having sequence No. 50, 51, 52, 53, 54 or 48, respectively) for a region of the highest score were used, and the first PCR was conducted with chromosomal DNA of AJ12092 strain as a template. The second PCR was conducted with the same chromosomal DNA as the template by using the PCR product as the primer for 3'-end and also using the primer 8 having sequence No. 49 as the primer on 5'-end to

15

obtain DNA fragments having the mutation introduced in the intended promoter region. To determine the activity of the mutant promoters, these DNA fragments were inserted into SmaI site of promoter probe vector pNEOL so that they were in the same direction with lacZ reporter gene to obtain plasmids pNEOL-1, pNEOL-2, pNEOL-3, pNEOL-4 and pNEOL-7. As a control for the activity, plasmid pNEOL-0 was constructed by inserting the DNA fragment, obtained by

20

[0127] pNEOL-0, pNEOL-1, pNEOL-2, pNEOL-3, pNEOL-4 and pNEOL-7 were introduced into AJ12092 strain, respectively. The plasmids were introduced by electrical pulse method (J. P. KOKAI No. 2-207791). The transformants were selected on CM2G medium plates (comprising 1 liter of pure water containing 10 g of polypeptone, 10 g of yeast extract, 5 g of glucose, 5 g of NaCl and 15 g of agar, and having pH 7.2) containing 4 μ g/ml of chloramphenicol, as chloramphenicol-resistant strains.

25

[0128] These strains were each spread on an agar medium (containing 0.5 g/dl of glucose, 1 g/dl of polypeptone, 1 g/dl of yeast extract, 0.5 g/dl of NaCl and 5 μ g/l of chloramphenicol), and cultured at 31.5°C for 20 hours. One aze of the cells thus obtained was inoculated into a medium [containing 3 g/dl of glucose, 1.5 g/dl of ammonium sulfate, 0.1 g/dl of KH₂PO₄, 0.04 g/dl of MgSO₄, 0.001 g/dl of FeSO₄, 0.01 g/dl of MnSO₄, 5 μ g/dl of VB₁, 5 μ g/dl of biotin and 45 mg/dl (in terms of N) of soybean hydrolyzate]. After the culture at 31.5°C for 18 hours, β -galactosidase activity of the

30

obtained cells was determined as described in Example 4(4). Since β -galactosidase activity was detected in AJ12092/pNEOL-0 as shown in Table 20, it was found that the DNA fragment inserted into the upstream of the gene of lacZ structure functioned as a promoter. In addition, β -galactosidase activity of each of the plasmid-introduced strains was higher than that of AJ12092/pNEOL-0. It was thus found that the transcription activity was increased by the introduction of the mutation into the promoter-like sequence, as shown in Table 20.

35

Table 20

	Relative activity (AJ12092/pNEOL-0=1)
AJ12092	nd
AJ12092/pNEOL-0	1.0
AJ12092/pNEOL-1	2.8
45 AJ12092/pNEOL-2	2.7
AJ12092/pNEOL-3	1.8
AJ12092/pNEOL-4	0.8
50 AJ12092/pNEOL-7	3.0

40

45

50

4) Construction of a plasmid for introduction of mutation:

- 55 [0129] PCR was conducted by using primers 14 and 15 (having the sequence of Seq ID No. 55 and Seq ID No.56) with chromosomal DNA of AJ12092 strain as the template. These DNA fragments thus obtained were inserted into a smaI site in a multicloning site of cloning vector pHSG398 (a product of TaKaRa) to construct plasmid p0. Then, p0 was digested with restriction enzymes EcoRV and BspHI, and also pNEOL-3 and pNEOL-7 were digested with restriction enzymes EcoRV and BspHI. DNA fragments thus obtained were ligated to obtain mutation-introducing plasmids p3

(mutant derived from mutation-introducing primer 11) and p7 (mutant derived from mutation-introducing primer 13).

5) Introduction of mutation-introducing plasmids into Arg-producing bacterium:

[0130] Each of the plasmids thus obtained was introduced into Arg-producing bacterium of the strain *Brevibacterium lactofermentum* AJ12092 by electrical pulse method (J. P. KOKAI No. 2-207791). Since these plasmids could not autonomously replicate in *Brevibacterium*, only the strains obtained by integrating these plasmids into the chromosome by homologous recombination could be selected as Cm-resistant strains. Strains in which the mutation-introducing plasmid was integrated into the chromosome were selected as chloramphenicol-resistant strains on CM2G medium plates (comprising 1 liter of pure water containing 10 g of polypeptone, 10 g of yeast extract, 5 g of glucose, 5 g of NaCl and 15 g of agar, and having pH 7.2) containing 5 µg/ml of chloramphenicol. Then, Cm-sensitive strains were selected in which the promoter region of *argG* gene was replaced with the intended modified sequence.

[0131] As a result, a strain substituted with P3 sequence (AJ12092-P3) and a strain substituted with P7 sequence (AJ12092-P7) were obtained.

(6) Cloning of *argG* gene

[0132] Based on the DNA sequence determined as in (1), oligonucleotides (primers 5 and 6) having the DNA sequence set out in Seq ID No. 46 and Seq ID No.47 were synthesized to conduct PCR using chromosomal DNA of *Brevibacterium flavum* as a template. The PCR reaction was conducted in 25 cycles, each cycle consisting of 94°C for 30 seconds, 55°C for one second and 72°C for 2 minutes and 30 seconds. The thus-obtained DNA fragment was cloned to SmaI site in multi-cloning site of cloning vector pSTV29 (Takara shuzo Co. Ltd.) to obtain pSTVargG. Furthermore, pargG was prepared by inserting into Sall site of pSTVargG a fragment containing the replication origin obtained by treating pSAK4 set out in Example 1 with Sall.

7) Introduction of pargG into Brev.:

[0133] pargG was introduced into the strain *Brevibacterium lactofermentum* AJ12092. Plasmid was introduced by electrical pulse method (J. P. KOKAI No. 2-207791). The transformant was selected as chloramphenicol-resistant strain on CM2G medium plates (comprising 1 liter of pure water containing 10 g of polypeptone, 10 g of yeast extract, 5 g of glucose, 5 g of NaCl and 15 g of agar, and having pH 7.2) containing 4 µg/ml of chloramphenicol.

8) ArgG activity of promoter modified strains:

[0134] ArgG activities of the above-described two kinds of *argG* promoter modified strains and a strain obtained by amplifying *argG* with plasmid (AJ12092/pargG) were determined. These strains were each spread on a agar medium (containing 0.5 g/dl of glucose, 1 g/dl of polypeptone, 1 g/dl of yeast extract, 0.5 g/dl of NaCl and 5 µg/l of chloramphenicol), and cultured at 31.5°C for 20 hours. One aze of the cells thus obtained were inoculated into a medium [containing 3 g/dl of glucose, 1.5 g/dl of ammonium sulfate, 0.1 g/dl of KH₂PO₄, 0.04 g/dl of MgSO₄, 0.001 g/dl of FeSO₄, 0.01 g/dl of MnSO₄, 5 µg/dl of VB₁, 5 µg/dl of biotin and 45 mg/dl (in terms of N) of soybean hydrolyzate]. After the culture at 31.5°C for 18 hours, ArgG activity of the obtained cells was determined by the method described above [Journal of General Microbiology (1990), 136, 1177-1183]. ArgG activities of the above-described two kinds of ArgG promoter modified strains and the strain (AJ12092/pargG) obtained by amplifying *argG* with plasmid are shown in Table 21. It is apparent from Table 21 that by introducing the mutation into the promoter, ArgG activity of AJ12092-P3 was increased to about twice as high as that of the parent strain, and the activity of AJ12092-P7 was increased to about three times as high as that of the parent strain. ArgG activity of AJ12092/pargG was about 4.5 times as high as that of the parent strain.

Table 21

	Relative activity (AJ12092=1)
AJ12092	1.0
AJ12092-P3	2.1
AJ12092-P7	2.9

Table 21 (continued)

	Relative activity (AJ12092=1)
AJ12092/pargG	4.4

9) Arg production by promoter modified strains:

[0135] The flask culture of each of argG promoter modified strains was conducted. As controls, parent strain AJ12092 and AJ12092/pargG were also cultured. These strains were each inoculated into a medium [containing 0.1 g/dl of KH_2PO_4 , 0.04 g/dl of MgSO_4 , 0.001 g/dl of FeSO_4 , 0.01 g/dl of MnSO_4 , 5 μ g/dl of VB_1 , 5 μ g/dl of biotin and 45 mg/dl (in terms of N) of soybean hydrolyzate]; and then spread on an agar medium (containing 0.5 g/dl of glucose, 1 g/dl of polypeptone, 1 g/dl of yeast extract, 0.5 g/dl of NaCl and 5 μ g/l of chloramphenicol), and cultured at 31.5°C for 20 hours. One aze of the cells were cultured in a flask containing 4 g/dl of glucose and 6.5 g/dl of ammonium sulfate at 31.5°C until glucose had been completely consumed. The absorbance (CD620) of the culture liquid diluted to a concentration of 1/51 with 0.2 N HCl solution, the quantity of arginine produced (concentration: g/dl) and culture time were shown in Table 22.

[0136] It is apparent from Table 22 that when argG promoter modified strain was used, the yield of arg was increased to a level equal to that of argG amplified with plasmid. As for the promoter varied strains, both AJ12092-P3 and AJ12092-P7 had the culture time equal to that of the parent strain, while the culture time of the plasmid amplified strain was increased. It was thus apparent that Arg productivity thereof was higher than that of the plasmid amplified strain.

Table 22

	OD	Arg (g/dl)	Culture time (h)	Productivity (g/dl/h)
AJ12092	0.502	1.25	48	0.026
AJ12092-P3	0.510	1.47	48	0.031
AJ12092-P7	0.514	1.43	48	0.030
AJ12092/pargG	0.520	1.47	52	0.028

Example 7 Introduction of mutation into GDH gene promoter region of coryneform glutamate-producing bacterium:

(1) Construction of mutant gdh plasmids:

[0137] Plasmids having GDH promoter sequence of FGR1 strain and FGR2 strain described in Example 2 were constructed by site directed mutagenesis. For obtaining GDH promoter sequence of FGR1 strain, PCR was conducted by using synthetic DNA shown in Seq ID No. 57 and synthetic DNA shown in No. 60 as the primers and chromosomal DNA of ATCC13869 as the template; and on the other hand, PCR was conducted by using synthetic DNA shown in Seq ID No. 58 and synthetic DNA shown in Seq ID No. 59 as the primers with chromosomal DNA of ATCC13869 as the template. Further, PCR was conducted by using synthetic DNAs shown in Seq ID Nos. 57 and Seq ID No.58 as the primers with a mixture of these PCR products as the template. The PCR product thus obtained was inserted into SmaI site of pSFKT2 (Japanese Patent Application No. 11-69896) to construct pSFKTG11. To obtain GDH promoter sequence of FGR2 strain, PCR was conducted by using synthetic DNA shown in Seq ID No. 57 and synthetic DNA shown in Seq ID No. 62 as the primers and chromosomal DNA of ATCC13869 as the template; and on the other hand, PCR was conducted by using synthetic DNA shown in Seq ID No. 58 and synthetic DNA shown in Seq ID No. 61 as the primers and chromosomal DNA of ATCC13869 as the template. Further, PCR was conducted by using synthetic DNA shown in Seq ID No. 57 and Seq ID No.58 as the primers and a mixture of these PCR products as the template. The PCR product thus obtained was inserted into SmaI site of pSFKT2 (Japanese Patent Application No. 11-69896) to construct pSFKTG07. The DNA sequences of the fragments inserted into SmaI sites of pSFKTG11 and pSFKTG07 were determined to confirm that no mutation was introduced into other regions than the promoter region in GDH.

(2) Construction of *gdh* promoter modified strains:

[0138] Then, pSFKTG11 and pSFKTG07 were introduced into AJ13029 strain by electrical pulse method, and transformants which grew on CM2B plates containing 25 μ g/ml of kanamycin at 25°C were selected. The transformants were cultured at 34°C to select strains which were resistant to kanamycin at 34°C. The fact that a strain is resistant to kanamycin at 34°C indicates that pSFKTG11 or pSFKTG07 was thus integrated on the chromosome of AJ13029 strain. Kanamycin-sensitive strains were obtained from the strains in which the plasmid was integrated on the chromosome. The GDH promoter sequences of these strains were determined. The strains having the same *gdh* promoter sequence as those of pSFKTG11 and pSFKTG07 were named GA01 and GA02, respectively.

(3) Confirmation of L-glutamic acid-productivity of *gdh* promoter modified strains:

[0139] The glutamic acid productivities of strains GA01 and GA02 and the parent strain AJ13029 were confirmed in the same manner as that of Example 2 (2) given above. As a result, a remarkable improvement in the accumulation of glutamic acid was recognized in GA01 and GA02 as shown in Table 23.

Table 23

Strain	Glu (g/dl)	Specific activity of GDH	Relative value
AJ13029	2.6	7.7	1.0
GA01	3.0	22.3	2.9
GA02	2.9	27.0	3.5

(4) Construction of self-cloning type *gdh* plasmid:

[0140] First, self-cloning vector pAJ220 was constructed. pAJ226 (J. P. KOKAI No.61-152289) was treated with EcoRV and PstI to prepare a fragment containing a region which could be autonomously replicated in a coryneform bacterium. The fragment was ligated with about 0.7 kb of the DNA fragment obtained by treating pAJ224 (J. P. KOKAI No. Sho 61-152289) with EcoRV and PstI to obtain a plasmid pAJ220. This plasmid could autonomously replicate in a coryneform bacterium, and it could afford trimethoprim resistance to the host.

[0141] PCR reaction was conducted by using synthetic DNA shown in Seq ID No. 63 and Seq ID No.64 as the primers and chromosomal DNA of wild-type coryneform bacterium strain ATCC13869 as the template. The *gdh* gene fragment thus obtained was inserted in Ball site of pAJ220 to construct pAJ220G. The promoter was present near Ball site of pAJ220, and the expression of the inserted gene was increased depending on the direction of the gene inserted into Ball site. pAJ220G and pGDH were introduced into ATCC13869 strain by electrical pulse method. GDH activities of the strains thus constructed were determined by the method stated in above-described step (1). As a result GDH activity of the strain into which pAJ220G had been introduced was about 1.5 times as high as that of the strain into which dGDH had been introduced as shown in Table 24.

Table 24

Strain	Specific activity of GDH	Relative value
ATCC13869	7.7	1.0
ATCC13869/pGDH	82.7	10.7
ATCC13869/pAJ220G	120.1	15.6

(5) Investigations on influence of *gdh* activity on the yield and by-produced Asp:

[0142] pGDH and pAJ220G were introduced into AJ13029 by electrical pulse method. Each of these strains and those obtained in above-described step (2) was inoculated into a seed culture medium having a composition shown in Table 25, and the shaking culture was conducted at 31.5°C for 24 hours to obtain the seed culture. 300 ml of medium for main culture having a composition shown in Table 25 was placed in each of 500 ml glass jar fermenters and then sterilized by heating. 40 ml of the seed cultures as described above were inoculated into the medium. The culture was started at a temperature of 31.5°C while the stirring rate and the aeration rate were controlled at 800 to 1300 rpm and

1/2 to 1/1 vvm, respectively. The pH of the culture liquid was kept at 7.5 with gaseous ammonia. The temperature was shifted to 37°C 8 hours after the initiation of the culture. The culture was terminated when glucose had been completely consumed in 20 to 40 hours, and the quantity of L-glutamic acid produced and accumulated in the culture liquid were determined (Table 26). The GDH activity for obtaining the highest yield was about 3-times as high. When GDH activity was further elevated, the degree of the improvement in the yield was reduced. When the GDH activity was elevated to about 16-times, the yield was rather reduced. Amino acids produced as by-products were analyzed with Hitachi Amino Acid Analyzer L-8500 to find that as GDH activity was elevated, the amount of accumulated aspartic acid and alanine was increased. These results proved the following facts: For increasing the yield of glutamic acid, it is necessary to suitably increase GDH activity so as not to cause a remarkable increase in the amount of aspartic acid and alanine. One of the effective methods therefor comprises the introduction of various mutations into gdh promoter to control GDH activity to about 3-times as high as that of the parent strain.

Table 25

Ingredient	Concentration	
	Seed culture	Main culture
Glucose	50 g/l	150 g/l
KH ₂ PO ₄	1 g/l	2 g/l
MgSO ₄ · 7H ₂ O	0.4 g/l	1.5 g/l
FeSO ₄ · 7H ₂ O	10 mg/l	15 mg/l
MnSO ₄ · 4H ₂ O	10 mg/l	15 mg/l
Soybean protein hydrolyzate	20 ml/l	50 ml/l
Biotin	0.5 mg/l	2 mg/l
Thiamine hydrochloride	2 mg/l	3 mg/l

Table 26

Strain	Glu (g/dl)	Asp (mg/dl)	Ala (mg/dl)	Relative activity of GDH	Relative value
AJ 13029	8.3	49	60	7.7	1.0
GA01	9.0	145	152	22.3	2.9
GA02	8.9	153	155	27.0	3.5
AJ13029/pGDH	8.8	201	190	82.7	10.7
AJ13029/pAJ220G	7.5	290	590	120.12	15.8

Sequence Listing

5

<110> Ajinomoto Co. Inc.

10

<120> Method of constructing amino acid producing bacteria, and method of preparing amino acids by fermentation with the constructed amino acid producing bacteria

15

<130> OP 99052

20

<150> JP 10-271786

<151> 1998-9-25

25

<150> JP 10-271787

<151> 1998-9-25

<160> 6

30

<210> 1

<211> 46

35

<212> nucleic acid

<400> 1

40

ttaattcttt gtggtcatat ctgcgacact gccataattt gaacgt

<210> 2

<211> 46

45

<212> nucleic acid

<400> 2

50

ttaattcttt gcggtcatat ctgcgacact gccataattt gaacgt

<210> 3

55

<211> 46

<212> nucleic acid

5

<400> 3

ttaattcttt gtggtcatat ctgcgacact gctataattt gaacgt

10

<210> 4

<211> 46

<212> nucleic acid

15

<400> 4

ttaattcttt gttgacatat ctgcgacact gctataattt gaacgt

20

<210> 5

<211> 46

<212> nucleic acid

25

<400> 5

ttaattcttt gttgccatat ctgcgacact gctataattt gaacgt

30

<210> 6

<211> 46

<212> nucleic acid

35

<400> 6

ttaattcttt gttgtcatat ctgcgacact gctataattt gaacgt

40

<210> 7

<211> 30

<212> nucleic acid

45

<220> primer A for cloning of gltA from Brevibacterium lactofermentum

<400> 7

gtcgacaata gcctgaatct gttctggtcg

50

<210> 8

55

<211> 30

<212> nucleic acid

<220> primer B for cloning of gltA from Brevibacterium lactofermentum

<400> 8

aagcttatacg acgctccct cccacggt

<210> 9

<211> 20

<212> nucleic acid

<220> primer 1 for introducing a mutation of gltA promoter

<400> 9

atcggtataa cgtgttaacc

<210> 10

<211> 20

<212> nucleic acid

<220> primer 2 for introducing a mutation of gltA promoter

<400> 10

atcggtataa tgtgttaacc

<210> 11

<211> 40

<212> nucleic acid

<220> primer 4 for introducing a mutation of gltA promoter

<400> 11

gatttgacaa aaccgcattt atcggtataa tgtgttaacc

<210> 12

<211> 28

<212> nucleic acid

<220> gltA promoter sequence primer

<400> 12
agggatccgt ccagtctcag acagcatc

5

<210> 13

<211> 17

<212> nucleic acid

<220> universal primer M13RV

10

<400> 13

caggaaacag ctatgac

15

<210> 14

<211> 20

<212> nucleic acid

<220> primer A for cloning of ICDH

20

<400> 14

gaattcgctc ccggtgcagc

25

<210> 15

<211> 20

<212> nucleic acid

<220> primer B for cloning of ICDH

30

<400> 15

gatgcagaat tccttgtcgg

35

<210> 16

<211> 28

<212> nucleic acid

<220> primer 1 for introducing a mutation of ICD promoter

40

<400> 16

tggattgctg gctataatgg tgtcgtga

45

<210> 17

50

55

<211> 53

<212> nucleic acid

<220> primer 2 for introducing a mutation of ICD promoter

<400> 17

caaccacgt tcagttgaca actactggat tgctggctat aatggtgtcg tga

<210> 18

<211> 53

<212> nucleic acid

<220> primer 3 for introducing a mutation of ICD promoter

<400> 18

caaccacgt tcagttgact actactggat tgctggctaa agtgggtgtcg tga

<210> 19

<211> 28

<212> nucleic acid

<220> primer 4 for introducing a mutation of ICD promoter

<400> 19

ggctgaaact gctataatag gcgccagc

<210> 20

<211> 51

<212> nucleic acid

<220> primer 5 for introducing a mutation of ICD promoter

<400> 20

ggaaacacgg cgttgccatg cggggctgaa actgctataa taggcgccag c

<210> 21

<211> 51

<212> nucleic acid

<220> primer 6 for introducing a mutation of ICD promoter

<400> 21

ggaaacacgg cgttgacatg cggggctgaa actgctataa taggcgccag c

5

<210> 22

<211> 22

<212> nucleic acid

10

<220> ICD promoter sequence primer

<400> 22

15

gtgcgggtcc agatgatctt ag

<210> 23

<211> 20

20

<212> nucleic acid

<220> primer A for amplifying of nptII

<400> 23

25

gggatcccg atgaatgtca

<400> 24

30

<211> 23

<212> nucleic acid

<220> primer B for amplifying of nptII

35

<400> 24

gcccgggggtg ggcgaagaac tcc

40

<210> 25

<211> 23

<212> nucleic acid

45

<220> primer for amplifying of *Brevibacterium lactofermentum pdhA gene LA*

<400> 25

aci gti tci atg ggi cti ggi cc

50

<210> 26

55

<211> 23

<212> nucleic acid

5 <220> primer for amplifying of *Brevibacterium lactofermentum pdhA gene LA*

<400> 26

10 cct tci ccg tti agi gti gti cg

<210> 27

<211> 30

15 <212> nucleic acid

<220> primer for *in vitro* cloning of *Brevibacterium lactofermentum pdhA gene LA*

20

<400> 27

ttg cag tta acc acg aag gtc agg ttg tcc

25

<210> 28

<211> 30

<212> nucleic acid

30 <220> primer for *in vitro* cloning of *Brevibacterium lactofermentum pdhA gene LA*

<400> 28

35 tgg atg aga cca cgt gat tct ggc tcg tcc

<210> 29

<211> 30

40

<212> nucleic acid

<220> primer for *in vitro* cloning of *Brevibacterium lactofermentum pdhA gene LA*

45

<400> 29

aca gat cct gca cga agg cat caa cga ggc

50

<210> 30

<211> 30

55

<212> nucleic acid

<220> primer for *in vitro* cloning of *Brevibacterium lactofermentum pdhA gene*
LA

<400>

tca tcg ctg cgg gta cct cct acg cca ccc

<210> 31

<211> 2766

<212> nucleic acid

<213> *Brevibacterium lactofermentum* ATCC13869

<220> *Brevibacterium lactofermentum* ATCC13869 *pdhA gene* LA

<400> 31

atg gcc gat caa gca aaa ctt ggt ggt aag ccc tcg gat gac tct aac 48

Met Ala Asp Gln Ala Lys Leu Gly Gly Lys Pro Ser Asp Asp Ser Asn

1 5 10 15

ttc gcg atg atc cgc gat ggc gtg gca tct tat ttg aac gac tca gat 96

Phe Ala Met Ile Arg Asp Gly Val Ala Ser Tyr Leu Asn Asp Ser Asp

20 25 30

ccg gag gag acc aac gag tgg atg gat tca ctc gac gga tta ctc cag 144

Pro Glu Glu Thr Asn Glu Trp Met Asp Ser Leu Asp Gly Leu Leu Gln

35 40 45

gag tct tct cca gaa cgt gct cgt tac ctc atg ctt cgt ttg ctt gag 192

Glu Ser Ser Pro Glu Arg Ala Arg Tyr Leu Met Leu Arg Leu Leu Glu

50 55 60

cgt gca tct gca aag cgc gta tct ctt ccc cca atg acg tca acc gac 240

Arg Ala Ser Ala Lys Arg Val Ser Leu Pro Pro Met Thr Ser Thr Asp

65 70 75 80

tac gtc aac acc att cca acc tct atg gaa cct gaa ttc cca ggc gat 288

Tyr Val Asn Thr Ile Pro Thr Ser Met Glu Pro Glu Phe Pro Gly Asp
 85 90 95
 5
 gag gaa atg gag aag cgt tac cgt cgt tgg att cgc tgg aac gca gcc 336
 Glu Glu Met Glu Lys Arg Tyr Arg Arg Trp Ile Arg Trp Asn Ala Ala
 10 100 105 110
 atc atg gtt cac cgc gct cag cga cca ggc atc ggc gtc ggc gga cac 384
 Ile Met Val His Arg Ala Gln Arg Pro Gly Ile Gly Val Gly Gly His
 15 115 120 125
 att tcc act tac gca ggc gca gcc cct ctg tac gaa gtt ggc ttc aac 432
 20 Ile Ser Thr Tyr Ala Gly Ala Ala Pro Leu Tyr Glu Val Gly Phe Asn
 130 135 140
 cac ttc ttc cgc ggc aag gat cac cca ggc ggc ggc gac cag atc ttc 480
 25 His Phe Phe Arg Gly Lys Asp His Pro Gly Gly Gly Asp Gln Ile Phe
 145 150 155 160
 ttc cag ggc cac gca tca cca ggt atg tac gca cgt gca ttc atg gag 528
 30 Phe Gln Gly His Ala Ser Pro Gly Met Tyr Ala Arg Ala Phe Met Glu
 165 170 175
 ggt cgc ctt tct gaa gac gat ctc gat ggc ttc cgt cag gaa gtt tcc 576
 35 Gly Arg Leu Ser Glu Asp Asp Leu Asp Gly Phe Arg Gln Glu Val Ser
 180 185 190
 cgt gag cag ggt ggc att ccg tcc tac cct cac cca cac ggt atg aag 624
 40 Arg Glu Gln Gly Gly Ile Pro Ser Tyr Pro His Pro His Gly Met Lys
 195 200 205
 gac ttc tgg gag ttc cca act gtg tcc atg ggt ctt ggc cca atg gat 672
 45 Asp Phe Trp Glu Phe Pro Thr Val Ser Met Gly Leu Gly Pro Met Asp
 50 210 215 220
 gcc att tac cag gca cgt ttc aac cgc tac ctc gaa aac cgt ggc atc 720
 55

Ala Ile Tyr Gln Ala Arg Phe Asn Arg Tyr Leu Glu Asn Arg Gly Ile
 225 230 235 240
 5
 aag gac acc tct gac cag cac gtc tgg gcc ttc ctt ggc gac ggc gaa 768
 Lys Asp Thr Ser Asp Gln His Val Trp Ala Phe Leu Gly Asp Gly Glu
 10 245 250 255
 atg gac gag cca gaa tca cgt ggt ctc atc cag cag gct gca ctg aac 816
 Met Asp Glu Pro Glu Ser Arg Gly Leu Ile Gln Gln Ala Ala Leu Asn
 15 260 265 270
 aac ctg gac aac ctg acc ttc gtg gtt aac tgc aac ctg cag cgt ctc 864
 Asn Leu Asp Asn Leu Thr Phe Val Val Asn Cys Asn Leu Gln Arg Leu
 20 275 280 285
 gac gga cct gtc cgc ggt aac acc aag atc atc cag gaa ctc gag tcc 912
 Asp Gly Pro Val Arg Gly Asn Thr Lys Ile Ile Gln Glu Leu Glu Ser
 25 290 295 300
 ttc ttc cgt ggc gca ggc tgg tct gtg atc aag gtt gtt tgg ggt cgc 960
 Phe Phe Arg Gly Ala Gly Trp Ser Val Ile Lys Val Val Trp Gly Arg
 30 305 310 315 320
 gag tgg gat gaa ctt ctg gag aag gac cag gat ggt gca ctt gtt gag 1008
 Glu Trp Asp Glu Leu Leu Glu Lys Asp Gln Asp Gly Ala Leu Val Glu
 35 325 330 335
 atc atg aac aac acc tcc gat ggt gac tac cag acc ttc aag gct aac 1056
 Ile Met Asn Asn Thr Ser Asp Gly Asp Tyr Gln Thr Phe Lys Ala Asn
 40 340 345 350
 gac ggc gca tat gtt cgt gag cac ttc ttc gga cgt gac cca cgc acc 1104
 Asp Gly Ala Tyr Val Arg Glu His Phe Phe Gly Arg Asp Pro Arg Thr
 45 355 360 365
 gca aag ctc gtt gag aac atg acc gac gaa gaa atc tgg aag ctg cca 1152
 Ala Lys Leu Val Glu Asn Met Thr Asp Glu Glu Ile Trp Lys Leu Pro
 50
 55

EP 1 033 407 A1

	370	375	380	
5	cgt ggc ggc cac gat tac cgc aag gtt tac gca gcc tac aag cga gct 1200			
	Arg Gly Gly His Asp Tyr Arg Lys Val Tyr Ala Ala Tyr Lys Arg Ala			
	385	390	395	400
10	ctt gag acc aag gat cgc cca acc gtc atc ctt gct cac acc att aag 1248			
	Leu Glu Thr Lys Asp Arg Pro Thr Val Ile Leu Ala His Thr Ile Lys			
	405	410	415	
15	ggc tac gga ctc ggc cac aac ttc gaa ggc cgt aac gca acc cac cag 1296			
	Gly Tyr Gly Leu Gly His Asn Phe Glu Gly Arg Asn Ala Thr His Gln			
20	420	425	430	
	atg aag aag ctg acg ctt gat gat ctg aag ttg ttc cgc gac aag cag 1344			
	Met Lys Lys Leu Thr Leu Asp Asp Leu Lys Leu Phe Arg Asp Lys Gln			
25	435	440	445	
	ggc atc cca atc acc gat gag cag ctg gag aag gat cct tac ctt cct 1392			
30	Gly Ile Pro Ile Thr Asp Glu Gln Leu Glu Lys Asp Pro Tyr Leu Pro			
	450	455	460	-
	cct tac tac cac cca ggt gaa gac gct cct gaa atc aag tac atg aag 1440			
35	Pro Tyr Tyr His Pro Gly Glu Asp Ala Pro Glu Ile Lys Tyr Met Lys			
	465	470	475	480
40	gaa cgt cgc gca gcg ctc ggt ggc tac ctg cca gag cgt cgt gag aac 1488			
	Glu Arg Arg Ala Ala Leu Gly Gly Tyr Leu Pro Glu Arg Arg Glu Asn			
	485	490	495	
45	TAC GAT CCA ATT CAG GTT CCA CCA CTG GAT AAG CTT CGC TCT GTC CGT 1536			
	Tyr Asp Pro Ile Gln Val Pro Pro Leu Asp Lys Leu Arg Ser Val Arg			
	500	505	510	
50	aag ggc tcc ggc aag cag cag atc gct acc act atg gcg act gtt cgt 1584			
	Lys Gly Ser Gly Lys Gln Gln Ile Ala Thr Thr Met Ala Thr Val Arg			
55				

EP 1 033 407 A1

	515	520	525	
5	acc ttc aag gaa ctg atg cgc gat aag ggc ttg gct gat cgc ctt gtc 1632			
	Thr Phe Lys Glu Leu Met Arg Asp Lys Gly Leu Ala Asp Arg Leu Val			
	530	535	540	
10	cca atc att cct gat gag gca cgt acc ttc ggt ctt gac tct tgg ttc 1680			
	Pro Ile Ile Pro Asp Glu Ala Arg Thr Phe Gly Leu Asp Ser Trp Phe			
	545	550	555	560
15	cca acc ttg aag atc tac aac ccg cac ggt cag aac tac gtg cct gtt 1728			
	Pro Thr Leu Lys Ile Tyr Asn Pro His Gly Gln Asn Tyr Val Pro Val			
	565	570	575	
20	gac cac gac ctg atg ctc tcc tac cgt gag gca cct gaa gga cag atc 1776			
	Asp His Asp Leu Met Leu Ser Tyr Arg Glu Ala Pro Glu Gly Gln Ile			
25	580	585	590	
	ctg cac gaa ggc atc aac gag gct ggt tcc gtg gca tcg ttc atc gct 1824			
30	Leu His Glu Gly Ile Asn Glu Ala Gly Ser Val Ala Ser Phe Ile Ala			
	595	600	605	
	gcg ggt acc tcc tac gcc acc cac ggc aag gcc atg att ccg ctg tac 1872			
35	Ala Gly Thr Ser Tyr Ala Thr His Gly Lys Ala Met Ile Pro Leu Tyr			
	610	615	620	
40	atc ttc tac tcg atg ttc gga ttc cag cgc acc ggt gac tcc atc tgg 1920			
	Ile Phe Tyr Ser Met Phe Gly Phe Gln Arg Thr Gly Asp Ser Ile Trp			
	625	630	635	640
45	gca gca gcc gat cag atg gca cgt ggc ttc ctc ttg ggc gct acc gca 1968			
	Ala Ala Ala Asp Gln Met Ala Arg Gly Phe Leu Leu Gly Ala Thr Ala			
	645	650	655	
50	ggt cgc acc acc ctg acc ggt gaa ggc ctc cag cac atg gat gga cac 2016			
	Gly Arg Thr Thr Leu Thr Gly Glu Gly Leu Gln His Met Asp Gly His			
55				

	660	665	670
5	tcc cct gtc ttg gct tcc acc aac gag ggt gtc gag acc tac gac cca 2064		
	Ser Pro Val Leu Ala Ser Thr Asn Glu Gly Val Glu Thr Tyr Asp Pro		
	675	680	685
10	tcc ttt gcg tac gag atc gca cac ctg gtt cac cgt ggc atc gac cgc 2112		
	Ser Phe Ala Tyr Glu Ile Ala His Leu Val His Arg Gly Ile Asp Arg		
	690	695	700
15	atg tac ggc cca ggc aag ggt gaa gat gtt atc tac tac atc acc atc 2160		
	Met Tyr Gly Pro Gly Lys Gly Glu Asp Val Ile Tyr Tyr Ile Thr Ile		
20	705	710	715 720
	tac aac gag cca acc cca cag cca gct gag cca gaa gga ctg gac gta 2208		
	Tyr Asn Glu Pro Thr Pro Gln Pro Ala Glu Pro Glu Gly Leu Asp Val		
25	725	730	735
	gaa ggc ctg cac aag ggc atc tac ctc tac tcc cgc ggt gaa ggc acc 2256		
	Glu Gly Leu His Lys Gly Ile Tyr Leu Tyr Ser Arg Gly Glu Gly Thr		
30	740	745	750 -
	ggc cat gag gca aac atc ttg gct tcc ggt gtt ggt atg cag tgg gct 2304		
	Gly His Glu Ala Asn Ile Leu Ala Ser Gly Val Gly Met Gln Trp Ala		
35	755	760	765
	ctc aag gct gca tcc atc ctt gag gct gac tac gga gtt cgt gcc aac 2352		
	Leu Lys Ala Ala Ser Ile Leu Glu Ala Asp Tyr Gly Val Arg Ala Asn		
40	770	775	780
	att tac tcc gct act tct tgg gtt aac ttg gct cgc gat ggc gct gct 2400		
	Ile Tyr Ser Ala Thr Ser Trp Val Asn Leu Ala Arg Asp Gly Ala Ala		
45	785	790	795 800
	cgt aac aag gca cag ctg cgc aac cca ggt gca gat gct ggc gag gca 2448		
	Arg Asn Lys Ala Gln Leu Arg Asn Pro Gly Ala Asp Ala Gly Glu Ala		
50			
55			

5 805 810 815
 ttc gta acc acc cag ctg aag cag acc tcc ggc cca tac gtt gca gtg 2496
 Phe Val Thr Thr Gln Leu Lys Gln Thr Ser Gly Pro Tyr Val Ala Val
 820 825 830
 10
 tct gac ttc tcc act gat ctg cca aac cag atc cgt gaa tgg gtc cca 2544
 Ser Asp Phe Ser Thr Asp Leu Pro Asn Gln Ile Arg Glu Trp Val Pro
 835 840 845
 15
 ggc gac tac acc gtt ctc ggt gca gat ggc ttc ggt ttc tct gat acc 2592
 Gly Asp Tyr Thr Val Leu Gly Ala Asp Gly Phe Gly Phe Ser Asp Thr
 20 850 855 860
 cgc cca gct gct cgt cgc ttc ttc aac atc gac gct gag tcc att gtt 2640
 25 865 870 875 880
 Arg Pro Ala Ala Arg Arg Phe Phe Asn Ile Asp Ala Glu Ser Ile Val
 gtt gca gtg ctg aac tcc ctg gca cgc gaa ggc aag atc gac gtc tcc 2688
 30 885 890 895 -
 Val Ala Val Leu Asn Ser Leu Ala Arg Glu Gly Lys Ile Asp Val Ser
 gtt gct gct cag gct gct gag aag ttc aag ttg gat gat cct acg agt 2736
 35 900 905 910
 Val Ala Ala Gln Ala Ala Glu Lys Phe Lys Leu Asp Asp Pro Thr Ser
 40 920 925 930
 gtt tcc gta gat cca aac gct cct gag gaa 2766
 Val Ser Val Asp Pro Asn Ala Pro Glu Glu
 45
 <210> 32
 <211> 8556
 50 <212> nucleic acid
 <213> Brevibacterium lactofermentum ATCC13869
 55

<400> 32

	tcacgttacg gcgatcaaca ccgcaaccac tacgagaaga tctccaaacg agaccaagag	60
5	cgcttctaag cccgtctcat ttgacacgtg ccattctgtg aggatatggc aggtgctttt	120
	tcatgccact atcttggggg tctcggtatt agatcttctg ataaaaaccc gatagttttc	180
	ttgcgctaga cactaattac ggcaccgctt aagcatggtc gtgacacgta aaacctgact	240
10	taggccattt tgatgtgggt tagatcatat tgacgtcaat gaatgaagtg actaactccg	300
	ccgaatccac atcgtctaaa aggcctgggc gaccacgtaa agacgggcac gacgagaaga	360
	tcacgcacgc aactttacgg ctcatcgaca gcaatcgtcc cgtcacggtc aatgcagttg	420
	tcaaagaaag cggagtggca cgtgcagcgg ttatctgacg ctggcccagg ctagtggatc	480
15	tagtagcgga agcttttagat gccgggagag ctccagttga aatagatacc ccaggggaca	540
	tcaaagagac cttgattgat gggctgttta caaatcaggc gaaaaccact ggagtctcct	600
	atcctcgta gcgatttcgc aaacggctcg agttgggtgat gtcagatcaa gaattacagc	660
20	tcgcctaata gaattcacat gtgaagagac gtcgagaagc aaatattcgc gcgctgcaag	720
	tcgcgcaaga aaaaggccaa atccggggcg atctagacat cgaggcgtgc ctcgatgcaa	780
	tccttggggg gttttattac caatcggtcg cgcgtggagt aaatttcacc gaccaaggta	840
25	caacgcaacg atgcagagaa gccttgaggg tgatctggca tggaatggaa ccttaaatc	900
	aggttctgac gaggtgcgaa gcaagttgtc gcgcgccgca cctcagtatc cggatcaact	960
	taatttcgaa gtgctgggtt ttctcgcgca taccatgc gtaccgatgt gccatgagc	1020
	gaaaaacagg ccacgataag tttcttaaaa cttatcgtgg cctgcttcta ttttgtgcg	1080
30	ccctgacggg ctgcaaccgc cgacctgctg ggtgtaaacc agctgctctt ccagctgagc	1140
	taaaggcgcg cactgtcttt tctagaacca ccttgggtggc ctgcaaagca acgagtgaaa	1200
	tactaacaca caatctccac agacctaaaa tcgctgctca ggccgtggaa attagcgatt	1260
35	gttaaggctt cttgtttcca cgctggacga ggcaagaacc ttgccaatta ccgagacgtt	1320
	ccgccttggg ctgcacgaga cctgccagtt gtgctgattc agagataact ccaggagcca	1380
	gggctccttc tttaccaatg ccaggagtca acaccagat acgaccattc tcagcgaggg	1440
40	agcggatgga atccacaagt ccgtcgacga gatcgccgtc atcctcgcg caccagagca	1500
	gcacgacatc gcacagctcg tcggtttctt catcgagtag ttcctcaccg attgcatctt	1560
	cgatggactc gctgatcagc gtgtcggaat cttcatccca tccaatttct tgaacgatat	1620
	gacccgattg aatgccgagt agttgagcat aatcctgggc accttgcttg actgcgccc	1680
45	gagcgtcggc cactttaata atcctctcgc tgtgggcccc gatgtgtttt tcgattacat	1740
	ggattcaaca tgaaccgcg gggtattga tatatccgaa ttgcacatta ccgtccaacc	1800
	ggtactttga accaccttcc cctggaattt tttccttttc ctccccctt acgctcaaga	1860
	atcaatgaat tcaatcactg gccagcgatt aacttttcga gttttcagtc ttggatttcc	1920
50	acaattctct tcaaaataat ggtggctaga tttttcatca aacctcacc aaaaggacat	1980
	cagacctgta gttttatgcg attcgcgtca aacgtgagag aaacatcaca tctcacggga	2040

55

EP 1 033 407 A1

	aactaccgga taattctttg caaaactttg caaagggtaa tgaacatgca gctagtttcc	2100
	gtagaaatgt tctttaaaaa atccacaaca attgccagga agcacaccga ttgatggata	2160
5	cctgaaatcc cagtgagcgc accactcccc ttacgtcaca gtctgtaaaa caaatcttcg	2220
	gtgttgcgta tccttggtta taacttatgc gttgacccat tcgtgcactt cgggtgtgcca	2280
	caattaggta cgaccaagaa tgggaccggg aaaccgggac gtataaacga aataaaacat	2340
10	tccaacagga ggtgtggaa atg gcc gat caa gca aaa ctt ggt ggt aag ccc	2392
	Met Ala Asp Gln Ala Lys Leu Gly Gly Lys Pro	
	5 10	
15	tcg gat gac tct aac ttc gcg atg atc cgc gat ggc gtg gca tct tat	2440
	Ser Asp Asp Ser Asn Phe Ala Met Ile Arg Asp Gly Val Ala Ser Tyr	
	15 20 25	
20	ttg aac gac tca gat ccg gag gag acc aac gag tgg atg gat tca ctc	2488
	Leu Asn Asp Ser Asp Pro Glu Glu Thr Asn Glu Trp Met Asp Ser Leu	
	30 35 40	
25	gac gga tta ctc cag gag tct tct cca gaa cgt gct cgt tac ctc atg	2536
	Asp Gly Leu Leu Gln Glu Ser Ser Pro Glu Arg Ala Arg Tyr Leu Met	
30	45 50 55	
	ctt cgt ttg ctt gag cgt gca tct gca aag cgc gta tct ctt ccc cca	2584
	Leu Arg Leu Leu Glu Arg Ala Ser Ala Lys Arg Val Ser Leu Pro Pro	
35	60 65 70 75	
	atg acg tca acc gac tac gtc aac acc att cca acc tct atg gaa cct	2632
40	Met Thr Ser Thr Asp Tyr Val Asn Thr Ile Pro Thr Ser Met Glu Pro	
	80 85 90	
45	gaa ttc cca ggc gat gag gaa atg gag aag cgt tac cgt cgt tgg att	2680
	Glu Phe Pro Gly Asp Glu Glu Met Glu Lys Arg Tyr Arg Arg Trp Ile	
	95 100 105	
50	cgc tgg aac gca gcc atc atg gtt cac cgc gct cag cga cca ggc atc	2728
	Arg Trp Asn Ala Ala Ile Met Val His Arg Ala Gln Arg Pro Gly Ile	
	110 115 120	
55		

5	ggc gtc ggc gga cac att tcc act tac gca ggc gca gcc cct ctg tac Gly Val Gly Gly His Ile Ser Thr Tyr Ala Gly Ala Ala Pro Leu Tyr	2776
	125 130 135	
10	gaa gtt ggc ttc aac cac ttc ttc cgc ggc aag gat cac cca ggc ggc Glu Val Gly Phe Asn His Phe Phe Arg Gly Lys Asp His Pro Gly Gly	2824
	140 145 150 155	
15	ggc gac cag atc ttc ttc cag ggc cac gca tca cca ggt atg tac gca Gly Asp Gln Ile Phe Phe Gln Gly His Ala Ser Pro Gly Met Tyr Ala	2872
	160 165 170	
20	cgt gca ttc atg gag ggt cgc ctt tct gaa gac gat ctc gat ggc ttc Arg Ala Phe Met Glu Gly Arg Leu Ser Glu Asp Asp Leu Asp Gly Phe	2920
	175 180 185	
25	cgt cag gaa gtt tcc cgt gag cag ggt ggc att ccg tcc tac cct cac Arg Gln Glu Val Ser Arg Glu Gln Gly Gly Ile Pro Ser Tyr Pro His	2968
	190 195 200	
30	cca cac ggt atg aag gac ttc tgg gag ttc cca act gtg tcc atg ggt Pro His Gly Met Lys Asp Phe Trp Glu Phe Pro Thr Val Ser Met Gly	3016
35	205 210 215	
40	ctt ggc cca atg gat gcc att tac cag gca cgt ttc aac cgc tac ctc Leu Gly Pro Met Asp Ala Ile Tyr Gln Ala Arg Phe Asn Arg Tyr Leu	3064
	220 225 230 235	
45	gaa aac cgt ggc atc aag gac acc tct gac cag cac gtc tgg gcc ttc Glu Asn Arg Gly Ile Lys Asp Thr Ser Asp Gln His Val Trp Ala Phe	3112
	240 245 250	
50	ctt ggc gac ggc gaa atg gac gag cca gaa tca cgt ggt ctc atc cag Leu Gly Asp Gly Glu Met Asp Glu Pro Glu Ser Arg Gly Leu Ile Gln	3160
	255 260 265	
55		

EP 1 033 407 A1

	cag gct gca ctg aac aac ctg gac aac ctg acc ttc gtg gtt aac tgc	3208
	Gln Ala Ala Leu Asn Asn Leu Asp Asn Leu Thr Phe Val Val Asn Cys	
5	270 275 280	
	aac ctg cag cgt ctc gac gga cct gtc cgc ggt aac acc aag atc atc	3256
10	Asn Leu Gln Arg Leu Asp Gly Pro Val Arg Gly Asn Thr Lys Ile Ile	
	285 290 295	
	cag gaa ctc gag tcc ttc ttc cgt ggc gca ggc tgg tct gtg atc aag	3304
15	Gln Glu Leu Glu Ser Phe Phe Arg Gly Ala Gly Trp Ser Val Ile Lys	
	300 305 310 315	
	gtt gtt tgg ggt cgc gag tgg gat gaa ctt ctg gag aag gac cag gat	3352
20	Val Val Trp Gly Arg Glu Trp Asp Glu Leu Leu Glu Lys Asp Gln Asp	
	320 325 330	
	ggg gca ctt gtt gag atc atg aac aac acc tcc gat ggt gac tac cag	3400
25	Gly Ala Leu Val Glu Ile Met Asn Asn Thr Ser Asp Gly Asp Tyr Gln	
	335 340 345	
30	acc ttc aag gct aac gac ggc gca tat gtt cgt gag cac ttc ttc gga	3448
	Thr Phe Lys Ala Asn Asp Gly Ala Tyr Val Arg Glu His Phe Phe Gly	
35	350 355 360	
	cgt gac cca cgc acc gca aag ctc gtt gag aac atg acc gac gaa gaa	3496
40	Arg Asp Pro Arg Thr Ala Lys Leu Val Glu Asn Met Thr Asp Glu Glu	
	365 370 375	
	atc tgg aag ctg cca cgt ggc ggc cac gat tac cgc aag gtt tac gca	3544
45	Ile Trp Lys Leu Pro Arg Gly Gly His Asp Tyr Arg Lys Val Tyr Ala	
	380 385 390 395	
	gcc tac aag cga gct ctt gag acc aag gat cgc cca acc gtc atc ctt	3592
50	Ala Tyr Lys Arg Ala Leu Glu Thr Lys Asp Arg Pro Thr Val Ile Leu	
	400 405 410	
55		

	gct cac acc att aag ggc tac gga ctc ggc cac aac ttc gaa ggc cgt	3640
	Ala His Thr Ile Lys Gly Tyr Gly Leu Gly His Asn Phe Glu Gly Arg	
5	415 420 425	
	aac gca acc cac cag atg aag aag ctg acg ctt gat gat ctg aag ttg	3688
10	Asn Ala Thr His Gln Met Lys Lys Leu Thr Leu Asp Asp Leu Lys Leu	
	430 435 440	
	ttc cgc gac aag cag ggc atc cca atc acc gat gag cag ctg gag aag	3736
15	Phe Arg Asp Lys Gln Gly Ile Pro Ile Thr Asp Glu Gln Leu Glu Lys	
	445 450 455	
	gat cct tac ctt cct cct tac tac cac cca ggt gaa gac gct cct gaa	3784
20	Asp Pro Tyr Leu Pro Pro Tyr Tyr His Pro Gly Glu Asp Ala Pro Glu	
	460 465 470 475	
	atc aag tac atg aag gaa cgt cgc gca gcg ctc ggt ggc tac ctg cca	3832
25	Ile Lys Tyr Met Lys Glu Arg Arg Ala Ala Leu Gly Gly Tyr Leu Pro	
	480 485 490	
30		
	gag cgt cgt gag aac tac gat cca att cag gtt cca cca ctg gat aag	3880
	Glu Arg Arg Glu Asn Tyr Asp Pro Ile Gln Val Pro Pro Leu Asp Lys	
35	495 500 505	
	ctt cgc tct gtc cgt aag ggc tcc ggc aag cag cag atc gct acc act	3928
40	Leu Arg Ser Val Arg Lys Gly Ser Gly Lys Gln Gln Ile Ala Thr Thr	
	510 515 520	
	atg gcg act gtt cgt acc ttc aag gaa ctg atg cgc gat aag ggc ttg	3976
45	Met Ala Thr Val Arg Thr Phe Lys Glu Leu Met Arg Asp Lys Gly Leu	
	525 530 535	
	gct gat cgc ctt gtc cca atc att cct gat gag gca cgt acc ttc ggt	4024
50	Ala Asp Arg Leu Val Pro Ile Ile Pro Asp Glu Ala Arg Thr Phe Gly	
	540 545 550 555	
55		

	ctt gac tct tgg ttc cca acc ttg aag atc tac aac ccg cac ggt cag	4072
5	Leu Asp Ser Trp Phe Pro Thr Leu Lys Ile Tyr Asn Pro His Gly Gln	
	560 565 570	
10	aac tac gtg cct gtt gac cac gac ctg atg ctc tcc tac cgt gag gca	4120
	Asn Tyr Val Pro Val Asp His Asp Leu Met Leu Ser Tyr Arg Glu Ala	
	575 580 585	
15	cct gaa gga cag atc ctg cac gaa ggc atc aac gag gct ggt tcc gtg	4168
	Pro Glu Gly Gln Ile Leu His Glu Gly Ile Asn Glu Ala Gly Ser Val	
	590 595 600	
20	gca tcg ttc atc gct gcg ggt acc tcc tac gcc acc cac ggc aag gcc	4216
	Ala Ser Phe Ile Ala Ala Gly Thr Ser Tyr Ala Thr His Gly Lys Ala	
	605 610 615	
25	atg att ccg ctg tac atc ttc tac tcg atg ttc gga ttc cag cgc acc	4264
	Met Ile Pro Leu Tyr Ile Phe Tyr Ser Met Phe Gly Phe Gln Arg Thr	
30	620 625 630 635	
35	ggt gac tcc atc tgg gca gca gcc gat cag atg gca cgt ggc ttc ctc	4312
	Gly Asp Ser Ile Trp Ala Ala Ala Asp Gln Met Ala Arg Gly Phe Leu	
	640 645 650	
40	ttg ggc gct acc gca ggt cgc acc acc ctg acc ggt gaa ggc ctc cag	4360
	Leu Gly Ala Thr Ala Gly Arg Thr Thr Leu Thr Gly Glu Gly Leu Gln	
	655 660 665	
45	cac atg gat gga cac tcc cct gtc ttg gct tcc acc aac gag ggt gtc	4408
	His Met Asp Gly His Ser Pro Val Leu Ala Ser Thr Asn Glu Gly Val	
	670 675 680	
50	gag acc tac gac cca tcc ttt gcg tac gag atc gca cac ctg gtt cac	4456
	Glu Thr Tyr Asp Pro Ser Phe Ala Tyr Glu Ile Ala His Leu Val His	
	685 690 695	

55

EP 1 033 407 A1

	cgt ggc atc gac cgc atg tac ggc cca ggc aag ggt gaa gat gtt atc	4504
	Arg Gly Ile Asp Arg Met Tyr Gly Pro Gly Lys Gly Glu Asp Val Ile	
5	700 705 710 715	
	tac tac atc acc atc tac aac gag cca acc cca cag cca gct gag cca	4552
10	Tyr Tyr Ile Thr Ile Tyr Asn Glu Pro Thr Pro Gln Pro Ala Glu Pro	
	720 725 730	
	gaa gga ctg gac gta gaa ggc ctg cac aag ggc atc tac ctc tac tcc	4600
15	Glu Gly Leu Asp Val Glu Gly Leu His Lys Gly Ile Tyr Leu Tyr Ser	
	735 740 745	
	cgc ggt gaa ggc acc ggc cat gag gca aac atc ttg gct tcc ggt gtt	4648
20	Arg Gly Glu Gly Thr Gly His Glu Ala Asn Ile Leu Ala Ser Gly Val	
	750 755 760	
	ggt atg cag tgg gct ctc aag gct gca tcc atc ctt gag gct gac tac	4696
25	Gly Met Gln Trp Ala Leu Lys Ala Ala Ser Ile Leu Glu Ala Asp Tyr	
	765 770 775	
	gga gtt cgt gcc aac att tac tcc gct act tct tgg gtt aac ttg gct	4744
30	Gly Val Arg Ala Asn Ile Tyr Ser Ala Thr Ser Trp Val Asn Leu Ala	
	780 785 790 795	
	cgc gat ggc gct gct cgt aac aag gca cag ctg cgc aac cca ggt gca	4792
40	Arg Asp Gly Ala Ala Arg Asn Lys Ala Gln Leu Arg Asn Pro Gly Ala	
	800 805 810	
	gat gct ggc gag gca ttc gta acc acc cag ctg aag cag acc tcc ggc	4840
45	Asp Ala Gly Glu Ala Phe Val Thr Thr Gln Leu Lys Gln Thr Ser Gly	
	815 820 825	
	cca tac gtt gca gtg tct gac ttc tcc act gat ctg cca aac cag atc	4888
50	Pro Tyr Val Ala Val Ser Asp Phe Ser Thr Asp Leu Pro Asn Gln Ile	
	830 835 840	

55

	cgt gaa tgg gtc cca ggc gac tac acc gtt ctc ggt gca gat ggc ttc	4936
	Arg Glu Trp Val Pro Gly Asp Tyr Thr Val Leu Gly Ala Asp Gly Phe	
5	845 850 855	
	ggt ttc tct gat acc cgc cca gct gct cgt cgc ttc ttc aac atc gac	4984
10	Gly Phe Ser Asp Thr Arg Pro Ala Ala Arg Arg Phe Phe Asn Ile Asp	
	860 865 870 875	
	gct gag tcc att gtt gtt gca gtg ctg aac tcc ctg gca cgc gaa ggc	5032
15	Ala Glu Ser Ile Val Val Ala Val Leu Asn Ser Leu Ala Arg Glu Gly	
	880 885 890	
	aag atc gac gtc tcc gtt gct gct cag gct gct gag aag ttc aag ttg	5080
20	Lys Ile Asp Val Ser Val Ala Ala Gln Ala Ala Glu Lys Phe Lys Leu	
	895 900 905	
	gat gat cct acg agt gtt tcc gta gat cca aac gct cct gag gaa taaat	5130
25	Asp Asp Pro Thr Ser Val Ser Val Asp Pro Asn Ala Pro Glu Glu	
	910 915 920	
30		
	cacctcaagg gacagataaa tcccgcggcc agacgttagt ctggcggcgg gattcgtcgt	5190
	aaagcaagct ctttttagcc gagaaacgcc ttgtcagaca atgttgcgcc cttgatattg	5250
35	gcgaactcct gcagcaaacc gcgcacagtc aacttcgact tggtagcctg atctgcctgg	5310
	tagacaatct ggccttcatg catcatgatc aggcgattgc ccaggcgaat tgcctgttcc	5370
	atgtttgtgcg tgaccataag cgtagtcaga gttccatctg ccacgatcctt ttcgggtcaag	5430
	gtgggtcacaa gctctgcacg ctgtggatca agcgtgcggg tgtgtctcatc caacagcatg	5490
40	attttaggtt gagtaaaacc agccatcagc agggacaatg cctgacgctg accgccagag	5550
	agcaaacc aa ctttggcagt gagcctgttt tccagaccca gctcaaggcg ctcaagttcc	5610
	tgcttgaatt gctcacggcg cttcgaggtc agtgcaaagc ccaatccacg gcgcttgccg	5670
45	cgcagcaacg cgatggccag attctcttca atggtgagat tcggcgcggg gcctgccaaa	5730
	ggatcctgaa aaacgcggcc gatgtagcgg gcaacgttgt gctctgacat cttgtttacc	5790
	ttgttgccgt cgatggaaat ctgcgaggaa tcaacaagca aacggccaga aacagcgttg	5850
	agcaggggtgg atttaccgcg accgttagaa ccgatgacgg tgacaaaatc gccctcagcc	5910
50	atctcgagtt tgagctgctg caacgcggcg cgctcattca cagtgcgggg gaagaaggtt	5970
	ttggaaattc cgttgatgga taacatgtct taagcctcca ctgctactgg ttgcttaggc	6030
55		

	ttcgggtgcct tggagaactt cgcacgccac ctcggcgagca gcatggcgac aaccaccaag	6090
	atcgcagaaa ttgccttcat atcgttgggg tcaaggccaa cgcgcagtgc tgcgaaaatg	6150
5	atcaggcggg acgcgatggc accgacgatg acagccaaca cagccaacca cacgcgacgc	6210
	tgaccgaaga tggcctggcc ccaaaataac cgatgcgaga ccgatcacga tgaggccaat	6270
	acccatcgaa atatctgcga agccctggta ctgagcgatg agtgcaccgg caagaccaac	6330
10	agaaccattg gacagggaga tgggtaggat tttggtgaaa tccgttgaaa caccaaagga	6390
	ctgcaccatc ggcccggtgt cgcgggtgga tcgcagcgac agtccgatat cagtgttgag	6450
	gaaccagatg acgatgagtc ccaaaaatcc cactgcaacg gcgaggatcg ccgggcctgc	6510
	ccatgtgccg aggaggccgg cgtcgcgaag cgggggtgaag aggttatcgg tgcgcaacaa	6570
15	tggcacgttc gcgccacca tgatgcgcaa gttaaccgac cacaacgcaa tcatggtcaa	6630
	aatacctgcg agcaaaccat cgatcttgcc cttggtgtgc agcaaaccgg tgatcatgcc	6690
	agcgataaag ccagtaacga aaccagcggc agtagccata agaggaggcc agccagacat	6750
20	aagagctgtc gcagctgttg ccgcgccagt ggtcaggctg ccgtcaacgg tgaggtcggg	6810
	aaagttgagc acacggaacg tcaaatagac gcccaatgcg acaactccgt acaacaatcc	6870
	gaactcaaaa gcgccgatca tacgcgttcg gccttatcca aaatctcttg agggatctcc	6930
	acgccctggc gctctgctgc atcttctgtg atcacgtagg tgaactcagt tgcagtctcc	6990
25	acaggcatgg ttgctgggtc ttcgccgtcc tgcagaatac gcagagccat ctcgccagtc	7050
	tggcggccaa gctcgggtga atcgataccc aggggttgcca gtgcgccacc ctcaacagtg	7110
	ccggactcag caccgatcac agggatctgc tctgtctcag caacctgaac cagagaagaa	7170
30	ataccgaaa caaccatgtt gtcagttgga acgtagatga catcaacatc gccgagagct	7230
	tcaacagcct gctgaatctc gttcacggta gtgacagtct gagtattaac ggacagcccc	7290
	agtggctcag cagccttggg gacctcatcg acctgcacct gagagttgac ctcaccagac	7350
	gcgtagacga tgccgatgga ctttgctgca ggaaccagct gctgcaaaag ctccaactgc	7410
35	tgctcaatcg gtgcgatatc agaagtaccg gtgacgtttc cgccagggtc ttcattagaa	7470
	tccaccagct ctgccgacac tgcacggta actgcggta acaggactgg gatatcagtg	7530
	atattctgcg cagttgcctg tgctgctgga gttgcaacag ccaacacgag atccaaattg	7590
40	tcagaagcga actgctgaga aatagtcagt gcagtgcctt gctcgcctt agcgttttgc	7650
	tcatcaaagg tgacgtcaac gcctgcctct tcaaaagctt ccttgaaacc agtggtcgtc	7710
	gcatcaagtg caggggtgctg aacaagctgg ttgatgcaa ctcggtaaga gtcgccacct	7770
	gcagcatcag tggaggtgga gctgtcactg gaatcgcttg agcacgaagc caacgccaag	7830
45	gcgccaacag taaagatgct tgcgagtacc ttcgaacggg aagaaaacat agcacatctc	7890
	cttaaagtgt tattttcaaa aaggggcaga cagcgtcaac acatgtctcg gataaagaac	7950
	catatgtgaa atgtctcatg atttaaaacta cttgttctac cagtcatatg cgcaattccc	8010
50	cctggatatc ccgcaggaca tggacaaaat ggggtgatag cgggtgcacc aattcaatct	8070
	tttaaaggcc ctagacaccg cgatttcttt aatcgatcat taaagaggga tcctctcccc	8130
	taacaaacct ccaaagacta gagtggggaa cccatgaac gtttctctca ataaaccag	8190

tgactctgac cgcgaatata ttcaatcaga actcaccgag ctggttgcc aggggagact 8250
 cgatctagat acttaccaag acgtggttga taccgtttgg tctactgatg atctaggcga 8310
 5 gttgatgagg atccgtgccc gcttctggg agggccgag gtttcgcagc agcggcccca 8370
 gcagccgag caaccacatc agcggccgca acagcaaccg ccacagcatt atggacaacc 8430
 cggctacggc caatcacctc aatatccacc gcagcagcct ccgcataatc agcccggtta 8490
 10 ttaccccgat cccggccctg gccagcagca accaccgatg caccagccac caacgcgtcc 8550
 aaatca

<210> 33
 15 <211> 20
 <212> nucleic acid
 <220> primer for construction of Brevibacterium lactofermentum pdhA gene LA
 20 amplification plasmid

<400> 33
 25 aat gcc agg agt caa cac cc

<210> 34
 <211> 20
 <212> nucleic acid
 30 <220> primer for construction of Brevibacterium lactofermentum pdhA gene LA
 amplification plasmid

35 <400> 34
 aca tgg aac agg caa ttc gc

40 <210> 35
 <211> 28
 <212> nucleic acid
 <220> primer for introducing a mutation of Brevibacterium lactofermentum pdhA
 45 gene LA promoter

<400> 35
 50 cgt ccc ggg ctg taa aac aaa tct tcg g

<210> 36
 55

<211> 27

<212> nucleic acid

5 <220> primer for introducing a mutation of Brevibacterium lactofermentum pdhA
gene LA promoter

<400> 36

10 atc ccc ggg ctt acc acc aag ttt tgc

<210> 37

15 <211> 30

<212> nucleic acid

<220> primer for introducing a mutation of Brevibacterium lactofermentum pdhA
gene LA promoter

20

<400> 37

ctt atg cgt tgc cac att cgt gca ctt cgg

25

<210> 38

<211> 40

<212> nucleic acid

30 <220> primer for introducing a mutation of Brevibacterium lactofermentum pdhA
gene LA promoter

<400> 38

35 gcg ttg acc cat tcg tgc act tcg gtg tgc tat aat tag g

<210> 39

<211> 40

40

<212> nucleic acid

<220> primer for introducing a mutation of Brevibacterium lactofermentum pdhA
gene LA promoter

45

<400> 39

gcg ttg cca cat tcg tgc act tcg gtg tgc tat aat tag g

50

<210> 40

55

<211> 38

<212> nucleic acid

5 <220> primer for introducing a mutation of Brevibacterium lactofermentum pdhA
gene LA promoter

<400> 40

10 ttt taa aac gtt ctg gag aag act cct gga gta atc cg

<210> 41

15 <211> 20

<212> nucleic acid

<220> primer for introducing a mutation of Brevibacterium lactofermentum pdhA
gene LA promoter

<400> 41

25 cga tct tgc ctt cgc gtg cc

<210> 42

<211> 30

30 <212> nucleic acid

<400> 42

35 agaccgccgg agtatgcaag aacgatgcgg

<210> 43

<211> 30

40 <212> nucleic acid

<400> 43

45 gacttcacca tcaatcatct tcttcaggta

<210> 44

<211> 30

50 <212> nucleic acid

<400> 44

55

accttcgacc agaccctggc taagggttt

5

<210> 45

<211> 30

<212> nucleic acid

10

<400> 45

gctaacaagc gcgatcgca agctggcaac

15

<210> 46

<211> 25

<212> nucleic acid

20

<400> 46

gcgatgacac cgtttttgtt ctgc

25

<210> 47

<211> 25

<212> nucleic acid

30

<400> 47

ggcgacatcc ttgccagat gatca

35

<210> 48

<211> 25

<212> nucleic acid

40

<400> 48

gattcacca tcaatcatct tcttc

45

<210> 49

<211> 24

<212> nucleic acid

50

<400> 49

gccaggtaca actgtctgaa ttgc

55

5
 <210> 50
 <211> 40
 <212> nucleic acid
 <220> primer for introducing a mutation

10
 <400> 50
 gttaatcgct tgccaatgca ggcaggtaag gtataacccg

15
 <210> 51
 <211> 40
 <212> nucleic acid

20
 <400> 51
 gttaatcgct tgctaattgca ggcaggtaag gtataacccg

25
 <210> 52
 <211> 40
 <212> nucleic acid

30
 <400> 52
 gttaatcgct tgtcaatgca ggcaggtaag gtataacccg

35
 <210> 53
 <211> 40
 <212> nucleic acid

40
 <400> 53
 gttaatcgct tgttaatgca ggcaggtaag gtataatccg

45
 <210> 54
 <211> 40
 <212> nucleic acid

50
 <400> 54
 gttaatcgct tgtcaatgca ggcaggtaag gtataatccg

55

5
 <210> 55
 <211> 30
 <212> nucleic acid

10
 <400> 55
 gggttcacagc ctcgtgcgga attcgtggag

15
 <210> 56
 <211> 25
 <212> nucleic acid

20
 <400> 56
 gcggttaccca gagctggatc ctcgg

25
 <210> 57
 <211> 16
 <212> nucleic acid

30
 <400> 57
 cagttgtggc tgatcg

35
 <210> 58
 <211> 17
 <212> nucleic acid

40
 <400> 58
 ctttccaga ctctggc

45
 <210> 59
 <211> 21
 <212> nucleic acid

50
 <400> 59
 gctataattt gacgtgagca t

55

<210> 60

<211> 25

5

<212> nucleic acid

<400> 60

10

gctcacgtca aattatagca gtgtc

<210> 61

<211> 54

15

<212> nucleic acid

<400> 61

20

ttgttgtcat tctgtgcgac actgctataa tttgaacgtg agcagttaac agcc

<210> 62

25

<211> 63

<212> nucleic acid

<400> 62

30

gttaactgct cacgttcaaa ttatagcaft gtcgcacaga atgacaacaa agaattaaaa ttg

<210> 63

35

<211> 25

<212> nucleic acid

<400> 63

40

gctagcctcg ggagctctct aggag

<210> 64

45

<211> 25

<212> nucleic acid

<400> 64

50

gatctttccc agactctggc cacgc

55

Claims

1. A method of producing coryneform bacteria having an improved amino acid- or nucleic acid-productivity, which comprises the steps of introducing a mutation in a promoter sequence of amino acid- or nucleic acid-biosynthesizing genes on a chromosome of a coryneform bacterium to make it close to a consensus sequence or introducing a change in a promoter sequence of amino acid- or nucleic acid-biosynthesizing genes on a chromosome of a coryneform bacterium by gene recombination to make it close to a consensus sequence, to obtain a mutant of the coryneform amino acid- or nucleic acid-producing microorganism, culturing the mutant and selecting a mutant capable of producing the intended amino acid or nucleic acid in a large amount.
2. The method of claim 1, wherein the amino acid is selected from the group consisting of glutamic acid, lysine, arginine, serine, phenylalanine, proline and glutamine, and nucleic acid is selected from the group consisting of inosine, guanosine, adenosine and nucleotide.
3. The method of claim 1, wherein the amino acid is glutamic acid, and the promoter for the biosynthesizing gene is selected from the group consisting of a promoter for glutamate dehydrogenase (GDH) gene, a promoter for citrate synthase (CS) gene, a promoter for isocitrate synthase (ICDH) gene, a promoter for pyruvate dehydrogenase (PDH) gene and a promoter for aconitase (ACO)-producing gene.
4. The method of claim 3, wherein the promoter for glutamate dehydrogenase (GDH) gene has a DNA sequence selected from the group consisting of (i) at least one DNA sequence selected from the group consisting of CGGTCA, TTGTCA, TTGACA and TTGCCA in -35 region (ii) TATAAT sequence or the same TATAAT sequence but in which the base of ATAAT is replaced with another base in -10 region, and (iii) a combination of (i) and (ii), wherein the sequence does not inhibit the function of the promoter.
5. The method of claim 4, wherein the promoter for GDH has TGGTCA in -35 region, and TATAAT in -10 region; or TTGTCA in -35 region, and TATAAT in -10 region.
6. The method of claim 3, wherein the promoter for CS has (i) TTGACA sequence in -35 region, (ii) TATAAT sequence in -10 region, or (iii) a sequence of the combination of (i) and (ii), and the sequence does not inhibit the function of the promoter.
7. The method of claim 3, wherein at least one of first and second promoters for ICDH has (i) TTGCCA or TTGACA sequence in -35 region, (ii) TATAAT sequence in -10 region, or (iii) a sequence of the combination of (i) and (ii), and the sequence does not inhibit the function of the promoter.
8. The method of claim 3, wherein the promoter for PDH has (i) TTGCCA sequence in -35 region, (ii) TATAAT sequence in -10 region, or (iii) a sequence of the combination of (i) and (ii), and the sequence does not inhibit the function of the promoter.
9. The method of claim 1, wherein the amino acid is arginine, and the promoter for the biosynthesizing gene is a promoter for argininosuccinate synthase.
10. The method of claim 9, wherein the promoter for the argininosuccinate synthase has a DNA sequence selected from the group consisting of (i) at least one DNA sequence selected from the group consisting of TTGCCA, TTGCTA and TTGTCA in -35 region, (ii) TATAAT sequence or TATAAT sequence but in which the base of ATAAC is replaced with another base in -10 region, and (iii) a combination of (i) and (ii) and the sequence does not inhibit the promoter function.
11. The method of claim 10, wherein the promoter for the argininosuccinate synthase has a DNA sequence selected from the group consisting of (i) TTGTCA in -35 region, (ii) TATAAT sequence in -10 region, and (iii) a combination of (i) and (ii).
12. A glutamic acid-synthesizing gene having a promoter according to any one of claims 4 to 8.
13. An arginine synthetase gene having the promoter of claim 10.
14. A coryneform glutamic acid-producing bacterium having the glutamate synthetase gene of claim 12.

15. A coryneform arginine-producing bacterium having the arginine synthetase gene of claim 13.

16. A method of producing an amino acid or nucleic acid by the fermentation, which comprises the steps of culturing a coryneform bacterium constructed by the method of any one claims 1 to 11 and having an improved amino acid- or nucleic acid-productivity, or the coryneform bacterium of claims 14 or 15 in a culture medium to form and thereby to accumulate the intended amino acid or nucleic acid in the culture medium, and collecting it from the culture medium.

17. A method of producing L-glutamic acid by fermentation, which comprises the steps of culturing a coryneform L-glutamic acid-producing bacterium resistant to 4-fluoroglutamic acid in a liquid culture medium to produce and thereby to accumulate L-glutamic acid in the culture medium, and collecting it from the culture medium.

FIG. 1




FIG. 2





FIG. 3



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/05175

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl.⁷ C12N15/67, C12N15/52, C12P13/04, C12P13/14,
C12P19/38, C12N9/02, C12N9/10, C12N1/21

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int. Cl.⁷ C12N15/67, C12N15/52, C12P13/04, C12P13/14,
C12P19/38, C12N9/02, C12N9/10, C12N1/21

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
BIOSIS (DIALOG), WPI (DIALOG), JICST (JOIS)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	FOURNIER, B. et al. "Strength and regulation of the different promoters for chromosomal β -lactamases of <i>Klebsiella Oxytoca</i> ", Antimicrobial Agents and Chemotherapy (April, 1999), Vol. 43, No. 4, pages 850-855	1-16
P, A	JEFFKE, T. et al. "Mutational analysis of the cbb operon (CO ₂ assimilation) promoter of <i>Ralstonia eutropha</i> ", J. Bacteriol., (July, 1999), Vol. 181, No. 14, pages 4374-4380	1-16
X	JP, 6-502548, A (Orsan), 24 March, 1994 (24.03.94) & WO, 93/03158, A1 & FR, 2679921, A1 & FR, 2679922, A1 & EP, 551506, A1	17
A	JP, 63-214189, A (Asahi Chemical Industry Co., Ltd.), 06 September, 1988 (06.09.88), (Family: none)	1-16

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
17 December, 1999 (17.12.99)

Date of mailing of the international search report
28 December, 1999 (28.12.99)

Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/05175

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Inventions as set forth in claims 1 to 16 relate to a process for preparing a corynebacterium having as improved amino acid- or nucleic acid-productivity characterized by preparing mutants by mutating the promoter sequence of an amino acid or nucleic acid biosynthesis gene on the chromosome of the corynebacterium, a gene having the promoter, the producing bacterium and a process for producing an amino acid or nucleic acid by culturing the producing bacterium.

Invention as set forth in claim 17 relates to a process for producing L-glutamic acid by culturing a L-glutamic acid-producing corynebacterium which is tolerant to 4-fluoroglutamic acid.

Such being the case, the inventions as set forth in claims 1 to 16 and the invention as set forth in claim 17 have no principal part in common and thus cannot be considered as forming a single general inventive concept.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1992)